#### For Research Use Only

# PE Anti-Mouse IFN gamma (XMG1.2)

Catalog Number:PE-65153 5 Publications



**Basic Information** 

Catalog Number:

PE-65153

100ug, 0.2 mg/ml

Source:

Isotype:

IgG1, kappa

GenBank Accession Number:

BC119063 GeneID (NCBI):

15978

**UNIPROT ID:** P01580

Full Name: interferon gamma

Calculated MW:

17 kDa

**Purification Method:** 

Affinity purification

CloneNo.: XMG1.2

Excitation/Emission maxima

wavelengths:

496 nm, 565 nm / 578 nm

**Applications** 

**Tested Applications:** 

Cited Applications:

FC (Intra)

Species Specificity:

Mouse

mouse

FC (Intra)

**Cited Species:** 

## **Background Information**

Interferon gamma (IFN  $\,\gamma$  ), is a type II interferon that provides immunity against bacterial, viral and protozoan infections. In its active form, IFN  $\gamma$  is a glycosylated, non-covalently linked homodimer of 29-32 kDa subunits. It is produced by a number of immune cell types including natural killer cells, natural killer T cells, and effector lymphocyte T cells following antigenic and inflammatory triggers. The IFN  $\gamma$  dimer binds to its cognate receptor which has two subunits: IFN-  $\gamma$  R1 which is the ligand-binding chain (  $\alpha$  chain) and IFN-  $\gamma$  R2, the signal-transducing chain (  $\beta$  chain). Binding to the receptor activates the JAK/STAT pathway which in turn activates IFN  $\gamma$  responsive genes. While IFN  $\gamma$  can inhibit viral replication, it also works as an immune-modulator and immune-stimulator by increasing surface expression of class I MHC proteins. (PMID: 19268625; 10688427)

### **Notable Publications**

Author	Pubmed ID	Journal	Application
Ping Yang	36184646	Nat Commun	FC (Intra)
Li Wu	35344716	Phytomedicine	FC (Intra)
Shan Fu	37926366	Cell Mol Gastroenterol Hepatol	FC (Intra)


Storage

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

Phosphate based buffer with 0.09% sodium azide and 0.1% gelatin, pH 7.2.

## Selected Validation Data



1X10^6 unstimulated or PMA and Ionomycin stimulated (in the presence of Brefeldin A) C57BL/6 mouse splenocytes were surface stained with CoraLite® Plus 647 Anti-Mouse CD3 (CL647-65077, Clone: 17A2) and then fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). Cells were then stained with 0.5 ug PE Anti-Mouse IFN gamma (PE-65153, Clone: XMG1.2).