

For Research Use Only

Vinculin Monoclonal antibody

Catalog Number: 66305-1-Ig **183 Publications**



Basic Information

Catalog Number: 66305-1-Ig	GenBank Accession Number: BC039174	Purification Method: Protein G purification
Concentration: 1000 ug/ml	GeneID (NCBI): 7414	CloneNo.: 2B5A7
Source: Mouse	UNIPROT ID: P18206	Recommended Dilutions: WB 1:5000-1:50000 IHC 1:50-1:500 IF/ICC 1:50-1:500
Isotype: IgG1	Full Name: vinculin	
Immunogen Catalog Number: AG24946	Calculated MW: 1133 aa, 124 kDa	
	Observed MW: 117 kDa	

Applications

Tested Applications: WB, IHC, IF/ICC, FC (Intra), ELISA	Positive Controls: WB : HepG2 cells, pig heart tissue, U-251 cells, LNCaP cells, U-87 MG cells, C6 cells, HEK-293 cells, HeLa cells, HuH-7 cells, LO2 cells, HSC-T6 cells, NIH/3T3 cells IHC : human breast cancer tissue, human prostate cancer tissue, mouse heart tissue IF/ICC : HeLa cells,
Cited Applications: WB, IHC, IF, CoIP	
Species Specificity: human, mouse, rat, pig	
Cited Species: human, mouse, rat, pig, zebrafish, hamster	

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

Vinculin belongs to the vinculin/alpha-catenin family. It is an actin filament (F-actin)-binding protein which involved in cell-matrix adhesion and cell-cell adhesion. Vinculin regulates cell-surface E-cadherin expression and potentiates mechanosensing by the E-cadherin complex. It may also play important roles in cell morphology and locomotion. Vinculin is a 117-kDa, 1,066-amino-acid protein which is ubiquitously expressed. Its splice variant, metavinculin (124 kDa), is muscle-specific.

Notable Publications

Author	Pubmed ID	Journal	Application
Juan M Barajas	34599880	Am J Pathol	WB
Tingben Huang	36167235	Acta Biomater	IF
Yizhou Zhu	31507175	ACS Appl Mater Interfaces	

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol, pH7.3
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 66305-1-Ig (Vinculin antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 66305-1-Ig (Vinculin Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 66305-1-Ig (Vinculin Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (10% Formaldehyde) fixed HeLa cells using 66305-1-Ig(Vinculin antibody) at dilution of 1:200 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



1×10^6 HeLa cells were intracellularly stained with 0.25 ug Vinculin Monoclonal antibody (66305-1-Ig, Clone:2B5A7) and CoraLite@488-Conjugated Goat Anti-Mouse IgG(H+L) (SA00013-1)(red), or 0.25 ug Mouse IgG1 isotype control Mouse McAb (66360-1-Ig, Clone: 1F8D3) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).