For Research Use Only

Mouse IgG2b isotype control Monoclonal antibody

Catalog Number:66360-3-lg 6 Publications



Basic Information

Catalog Number: 66360-3-lg Size:

Source: Mouse Isotype: IgG2b

1000 µg/ml

GenBank Accession Number:

GeneID (NCBI):
Full Name:

Purification Method: Protein A purification

CloneNo.: 11B8C4

Recommended Dilutions: WB 1:500-1:2000 IHC 1:1500-1:6000

Applications

Tested Applications: WB, IHC, FC, ELISA Cited Applications: WB, chIP

Species Specificity: megathura crenulata

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB:

Background Information

This antibody is used as the isotype control of mouse IgG2b. It can be used as the isotype control in Flow Cytometry, Immuno-Precipitation and other experiments. The immunogen of this antibody is Keyhole Limpet Hemocyanin (KLH).

Notable Publications

Author	Pubmed ID	Journal	Application
Kai Liu	36186122	Aging Dis	WB
Liang Liu	34381702	Front Oncol	ChIP
Yan Yu	32332880	Cell Death Dis	chIP

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates (50 ug/lane) were subjected to SDS PAGE followed by western blot with 66360-3-Ig (Mouse IgG2b isotype control antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours



Immunohistochemical analysis of paraffinembedded human colon tissue slide using 66360-3-Ig (Mouse IgG2b isotype control antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human colon tissue slide using 66360-3-Ig (Mouse IgG2b isotype control antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1x10^6 human PBMCs were surface stained with 0.5 ug Mouse I gG2b isotype control Monoclonal antibody (66360-3-Ig, Clone: 11B8C4) and Coralite® 488-Conjugated Goat Anti-Mouse IgG(H+L) (SA00013-1) (red), or 0.5 ug Anti-Human CD20 (65085-1-Ig) and Coralite® 488-Conjugated Goat Anti-Mouse IgG(H+L) (SA00013-1) (green), or unstained (blue). Cells were not fixed.