For Research Use Only

CoraLite® Plus 488-conjugated PAX8 Polyclonal antibody



Catalog Number: CL488-10336

Featured Product

Basic Information

Catalog Number:

CL488-10336

Size: 1000 ug/ml

Source: Rabbit Isotype:

Immunogen Catalog Number:

AG0306

GenBank Accession Number:

BC001060

GeneID (NCBI):

UNIPROT ID: Q06710

Full Name: paired box 8

Calculated MW: 48 kDa

Purification Method: Antigen affinity purification Recommended Dilutions:

IF/ICC 1:50-1:500 Excitation/Emission maxima

wavelengths: 493 nm / 522 nm

Applications

Tested Applications: IF/ICC, FC (Intra) Species Specificity:

human, mouse, rat

Positive Controls:

IF/ICC: SKOV-3 cells,

Background Information

PAX8 is a member of the paired box (PAX) family of transcription factors, typically containing a paired box domain and a paired-type homeodomain. It is expressed during organogenesis of the thyroid gland, kidney and Mullerian system. It is thought to regulate the expression of Wilms tumor suppressor (WT1) gene and mutations in PAX8 have $been \, associated \, with \, Wilms \, tumor \, cells, thy roid \, and \, ovarian \, carcinomas. \, PAX8 \, is \, a \, useful \, marker \, in \, distinguishing$ ovarian carcinomas from mammary carcinomas (PMID: 18724243). PAX8 is expressed in a high percentage of ovarian serous, endometrioid, and clear cell carcinomas, but only rarely in primary ovarian mucinous adenocarcinomas.

This antibody can detect all 5 isoforms of PAX8 (31 kDa, 34 kDa, 41 kDa, 43 kDa and 48 kDa). A 58-60 kDa form has also been reported (PMID:15650356, 1049952). Proteintech's 10336-1-AP antibody has cross-reactivities with the other PAX family members.


Storage

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



1X10^6 SKOV-3 cells were intracellularly stained with 0.2 ug Coralite® Plus 488 Anti-Human PAX8 (CL488-10336) (red), or 0.2 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).



Immunofluorescent analysis of (4% PFA) fixed SKOV-3 cells using CoraLite® Plus 488 PAX8 antibody (CL488-10336) at dilution of 1:200, CL594-phalloidin (red).