

## Human CD3 Magnetic Beads

Catalog#/Size: MS002-10/10 test

MS002-100/100 test

---

### Description


CD3 is a multimeric protein associated with T cell receptor (TCR) to form a complex involved in antigen recognition and signal transduction. 45%-70% human peripheral blood mononuclear cells (PBMC) express CD3. Human CD3 Magnetic Beads are used for isolation or depletion of human CD3 T lymphocytes from PBMC, whole blood, or other sample types. Following incubation with human CD3 antibody conjugated magnetic beads, the cell sample is placed on a magnet. CD3+ cells remain attached to magnetic beads after separation and can be used for further downstream applications, such as in cell expansion, but not suitable for flow cytometry analysis. CD3- cells remain in supernatant and could also be used for further applications.

### Product Details

Components	MS002-10: 200 $\mu$ L 10mg/mL Human CD3 Magnetic Beads MS002-100: 2x1mL 10mg/mL Human CD3 Magnetic Beads
Reactivity	Human
Beads Diameter	2.7 $\mu$ m
Storage buffer	PBS, pH7.4, 0.2% BSA and 0.05% Sodium Azide
Storage temperature	2-8 $^{\circ}$ C
Recommend usage	20 $\mu$ L Human CD3 Magnetic Beads for $1 \times 10^7$ cells

## Representative example of depletion

Following depletion of CD3+ cells, supernatant cell suspension was stained with FITC-CD45(F10-89-4) and CL647-CD3(UCHT1) antibodies. Left panel: CD3+CD45+ cells before selection. Right panel: CD3+CD45+ cells after depletion. All viable cells are gated in the analysis. Human CD3 magnetic beads are tested using PBMC from three different donors.



For technical support for this product please contact:

T: 1(888)4PTGLAB(1-888-478-4522) (toll free in USA), or 1(312)455-8498 (outside USA)

E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)

W: [ptglab.com](http://ptglab.com)