For Research Use Only

## Progranulin/PGRN Polyclonal antibody, PBS Only




Catalog Number:10053-1-PBS


Basic Information	Catalog Number: 10053-1-PBS	GenBank Accession Number: BC010577	Purification Method: Antigen affinity purification
	Size: 1 mg/ml	GenelD (NCBI): 2896	
	Source: Rabbit	UNIPROT ID: P28799	
	lsotype: IgG	Full Name: granulin	
	Immunogen Catalog Number: AG0010	Calculated MW: 64 kDa	
		Observed MW: 64 kDa	
Applications	Tested Applications: WB, IHC, Indirect ELISA		
	Species Specificity: human		
Background Information	GRN, also known as PGRN or PCDGF, is a cysteine-rich protein of 68.5 kDa that is typically secreted into a highly glycosylated 88 kDa form. PGRN is a unique growth factor that plays an important role in cutaneous wound healing. It has an anti-inflammatory effect and promotes cell proliferation. When PCDGF is degraded to several 6-25 kDa fragments, called granulins (GRNs) by neutrophil proteases, a pro-inflammatory reaction occurs. PGRN is widely expressed, particularly in epithelial cells, immune cells, neurons, and chondrocytes. High levels of PGRN expression have been reported in human cancers, and its expression is closely correlated with the development and metastasis of several cancers. The recent discovery that mutations in the gene encoding for pro-granulin (GRN) cause frontotemporal lobar degeneration (FTLD), and other neurodegenerative diseases leading to dementia, has brought renewed interest in progranulin and its functions in the central nervous system. Several in vitro studies have revealed that PGRN is classified into two isoforms according to its glycosylation status, the glycosylated immature isoform (58-68 kDa) and the fully glycosylated mature secretory isoform (88 kDa) (PMID: 25838514).		
Storage	Storage: Store at -80°C. The product is shipped with ice pa Storage Buffer: PBS Only	cks. Upon receipt, store it immediatel	yat-80°C

For technical support and original validation data for this product please contact: T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com


This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data





Various lysates were subjected to SDS PAGE followed by western blot with 10053-1-AP (Granulin antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 10053-1-PBS in a different storage buffer formulation. Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 10053-1-AP (Granulin antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 10053-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 10053-1-AP (Granulin antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 10053-1-PBS in a different storage buffer formulation.