### For Research Use Only

# PREB Polyclonal antibody

Catalog Number: 10146-2-AP

Featured Product

8 Publications



**Basic Information** 

Catalog Number:

10146-2-AP

Size:

500 ug/ml

Source:

Rabbit

Sotype:

GenBank Accession Number:

BC002765

GeneID (NCBI):

10113

UNIPROT ID:

Q9HCU5

Full Name:

G prolactin regulatory element binding

Immunogen Catalog Number: Calculated MW:

TE buffer pH 9.0; (\*) Alternatively, antigen

AG0202 45 kDa

Observed MW: 45 kDa

**Applications** 

Tested Applications: WB, IHC, IF/ICC, IP, ELISA

Cited Applications: WB, IF, IP, CoIP Species Specificity: human, mouse Cited Species:

human, mouse, rat

Note-IHC: suggested antigen retrieval with

retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: SKOV-3 cells, 3T3-L1 cells, HT-1080 cells, HepG2

**Purification Method:** 

WB 1:500-1:1000

IHC 1:500-1:2000

IF/ICC 1:50-1:500

protein lysate

Antigen affinity purification

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

cells

IP: SKOV-3 cells,

IHC: mouse intestine, human ovary tumor tissue,

human prostate cancer tissue

IF/ICC: HepG2 cells,

# **Background Information**

PREB contains two PQ-rich potential transactivation domains, but no apparent DNA-binding motif, and exhibits sequence-specific binding to Pit1-binding element of the prolactin (PRL) promoter. It may act as a transcriptional regulator and is thought to be involved in some of the developmental abnormalities observed in patients with partial trisomy 2p.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Mengyue You	36252341	Redox Biol	WB
Jiaxin Sun	28515233	J Cell Sci	WB,CoIP
Joo-Man Park	29601978	Biochim Biophys Acta	WB

Storage

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**




WB result of anti-PREB (10146-2-AP, 1:250) with HepG2 and HT1080 by Dr. Dzmitry Batrakou.




SKOV-3 cells were subjected to SDS PAGE followed by western blot with 10146-2-AP (PREB antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.




IP result of anti-PREB (IP:10146-2-AP, 4ug; Detection:10146-2-AP 1:500) with SKOV-3 cells lysate 3200ug.



Immunohistochemical analysis of paraffinembedded human prostate cancer tissue slide using 10146-2-AP (PREB Antibody) at dilution of 1:50 (under 40x lens).



Immunohistochemical analysis of paraffinembedded mouse intestine tissue slide using 10146-2-AP (PREB antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using PREB antibody (10146-2-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).