

For Research Use Only

DIDO1 Polyclonal antibody

Catalog Number: 10183-1-AP



Basic Information

Catalog Number: 10183-1-AP	GenBank Accession Number: BC000770	Purification Method: Antigen affinity purification
Size: 750 µg/ml	GeneID (NCBI): 11083	Recommended Dilutions: WB 1:1000-1:6000 IHC 1:50-1:500 IF/ICC 1:50-1:500
Source: Rabbit	UNIPROT ID: Q9BTC0	
Isotype: IgG	Full Name: death inducer-obliterator 1	
Immunogen Catalog Number: AG0237	Calculated MW: 59 kDa Observed MW: 300-350 kDa	

Applications

Tested Applications:
WB, IHC, IF/ICC, ELISA

Species Specificity:
human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: Raji cells,
IHC: human breast cancer tissue,
IF/ICC: HepG2 cells,

Background Information

DIDO1, also termed DATF-1, C20orf158, DATF1, KIAA0333 and DIO-1, is a putative transcription factor, weakly pro-apoptotic when overexpressed. It is a tumor suppressor. In mice, DIDO1 is upregulated by apoptotic signals and encodes a cytoplasmic protein that translocates to the nucleus upon apoptotic signal activation. When overexpressed, the mouse protein induced apoptosis in cell lines growing in vitro. The human DIDO1 is similar to the mouse gene and therefore is thought to be involved in apoptosis.

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Raji cells were subjected to SDS PAGE followed by western blot with 10183-1-AP (DIDO1 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 10183-1-AP (DIDO1 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 10183-1-AP (DIDO1 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using DIDO1 antibody (10183-1-AP) at dilution of 1:200 and Coralite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-phalloidin (red).