

## MMP7 Polyclonal antibody

Catalog Number: 10374-2-AP

Featured Product

97 Publications

## Basic Information

## Catalog Number:

10374-2-AP

## Source:

Rabbit

## Isotype:

IgG

## Immunogen Catalog Number:

AG0550

## GenBank Accession Number:

BC003635

## GeneID (NCBI):

4316

## UNIPROT ID:

P09237

## Full Name:

matrix metalloproteinase 7  
(matrilysin, uterine)

## Calculated MW:

29 kDa

## Observed MW:

28-30 kDa

## Purification Method:

Antigen affinity purification

## Recommended Dilutions:

WB: 1:1000-1:4000

IHC: 1:50-1:500

IF-P: 1:50-1:500

IF/ICC: 1:200-1:800

## Applications

## Tested Applications:

WB, IHC, IF/ICC, IF-P, ELISA

## Cited Applications:

WB, IHC, IF

## Species Specificity:

human, mouse

## Cited Species:

human, mouse, rat, pig

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

## Positive Controls:

**WB**: A549 cells, human placenta tissue, SKOV-3 cells, NIH/3T3 cells, COLO 320 cells, PC-3 cells

**IHC**: human pancreas cancer tissue, human prostate cancer tissue, human colon cancer tissue, human stomach cancer tissue

**IF-P**: human pancreas cancer tissue,

**IF/ICC**: PC-3 cells,

## Background Information

Matrix metalloproteinase-7 (MMP-7)/ matrilysin is a member of the MMP family, but is structurally different from the other MMPs by virtue of the absence of a conserved COOH-terminal protein domain. MMPs are involved in the breakdown of extracellular matrix in normal physiological processes, such as embryonic development reproduction, and tissue remodeling, as well as in disease processes, such as arthritis and cancer metastasis. Most MMP's are secreted as inactive proproteins which are activated when cleaved by extracellular proteinases. MMP-7 degrades proteoglycans, fibronectin, elastin and casein, and is involved in wound healing, tumor progression, pulmonary fibrosis, and development of choroidal neovascularization in age-related macular degeneration. The expression of MMP-7 is increased in most tumors. This antibody can only recognize the full-length of MMP7.

## Notable Publications

Author	Pubmed ID	Journal	Application
Rongfa Yuan	25056121	Cancer Res	WB
Xiao-Xiao He	30219235	Biochem Biophys Res Commun	WB
Xudong Peng	30205370	Cell Physiol Biochem	WB

## Storage

## Storage:

Store at -20°C. Stable for one year after shipment.

## Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926


E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)W: [ptgcn.com](http://ptgcn.com)

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 10374-2-AP (MMP7 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human pancreas cancer tissue slide using 10374-2-AP (MMP7 Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human pancreas cancer tissue slide using 10374-2-AP (MMP7 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed human pancreas cancer tissue using 10374-2-AP (MMP7 antibody) at dilution of 1:50 and Alexa Fluor 488-Conjugated Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (-20°C Methanol) fixed PC-3 cells using MMP7 antibody (10374-2-AP) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L).