For Research Use Only

Plexin B2 Polyclonal antibody

Catalog Number:10602-1-AP

Featured Product 9 Publications




Basic Information	Catalog Number: 10602-1-AP	GenBank Accessi BC004542	on Number:	Purification Method: Antigen affinity purification	
	Size: GeneID (NCBI): 1000 µ g/ml 23654 Source: UNIPROT ID:			Recommended Dilutions: WB 1:500-1:1000 IP 0.5-4.0 ug for 1.0-3.0 mg of total	
	Rabbit	015031		protein lysate IHC 1:400-1:1600 IF/ICC 1:10-1:100	
	Isotype: IgG Immunogen Catalog Number: AG0941	Full Name: plexin B2			
		Calculated MW: 205 kDa			
		Observed MW: 80 kDa, 240 kDa			
Applications	WB, IHC, IF/ICC, IP, ELISA Cited Applications: WB, IHC, IF, IP Species Specificity: human, mouse, rat Cited Species:			Positive Controls: WB : A549 cells, HEK-293 cells, HeLa cells, HepG2 cells, human brain tissue, MCF-7 cells, mouse liver tissue, mouse ovary tissue	
			human bra		
			IP: mouse	IP : mouse liver tissue,	
				IHC : human tonsillitis tissue, human liver cancer tissue, human breast cancer tissue, human gliomas tissue	
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0		IF/ICC : He	IF/ICC : HepG2 cells,	
Background Information	Semaphorins, originally identified as axon guidance molecules, have also been implicated in angiogenesis, immunoregulation, and cancer (PMID: 16584533). Plexins are main receptors for semaphorins and are thought to control many of the functional effects of semaphorins. PLXNB2 is one of the three plexin-B family members which participate in axon guidance and cell migration in response to semaphorins. It has been reported that plexin B2 precursor is 240 kDa and can be cleaved into alpha subunit (~170-kDa) and beta subunit (80-kDa) (PMID: 12533544). This antibody raised against 1542-1841aa of human plexin B2 can recognize precursor and beta subunit of plexin B2.				
Notable Publications	Author	Dubmed ID	lournal	Amilation	
	Katrin Gerstmann		Sci Adv	Application IF	
	Xiaonan Liu		Nat Commun	WB	
	Rongpan Bai		EMBO J	WB,IF	
Storage	Storage: Store at -20°C. Stable for one yea Storage Buffer: PBS with 0.02% sodium azide ar Aliquoting is unnecessary for -20	nd 50% glycerol pH 7.3.			

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data

for 1.5 hours.







Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 10602-1-AP (Plexin B2 antibody) at dilution of 1:800 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). A 549 cells were subjected to SDS PAGE followed by western blot with 10602-1-AP (Plexin B2 antibody) at dilution of 1:800 incubated at room temperature



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 10602-1-AP (Plexin B2 antibody) at dilution of 1:800 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-Plexin B2 (IP:10602-1-AP, 4ug; Detection:10602-1-AP 1:1000) with mouse liver tissue lysate 4000ug.



Immunofluorescent analysis of HepG2 cells using 10602-1-AP (Plexin B2 antibody) at dilution of 1:25 and Alexa Fluor 488-conjugated Goat Anti-Rabbit IgG(H+L).