

For Research Use Only

# P16-INK4A Polyclonal antibody

Catalog Number: 10883-1-AP

Featured Product

430 Publications



## Basic Information

### Catalog Number:

10883-1-AP

### Concentration:

650 ug/ml

### Source:

Rabbit

### Isotype:

IgG

### Immunogen Catalog Number:

AG1328

### GenBank Accession Number:

BC021998

### GeneID (NCBI):

1029

### UNIPROT ID:

P42771

### Full Name:

cyclin-dependent kinase inhibitor 2A

### Calculated MW:

16 kDa

### Observed MW:

16-18 kDa

### Purification Method:

Antigen affinity purification

### Recommended Dilutions:

WB: 1:1000-1:6000

IP: 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IHC: 1:1000-1:4000

IF/ICC: 1:200-1:800

FC (Intra): 0.40 ug per 10<sup>6</sup> cells in a 100 µl suspension

## Applications

### Tested Applications:

WB, IHC, FC (Intra), IP, ELISA

### Cited Applications:

WB, IHC, IP, CoIP

### Species Specificity:

human, dog

### Cited Species:

human, pig, rabbit, canine, monkey, bovine

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

### Positive Controls:

WB: HEK-293 cells, HEK293 cells, HeLa cells, HepG2 cells, PC-3 cells

IP: HEK-293 cells,

IHC: human cervical cancer tissue,

IF/ICC: MDCK cells,

FC (Intra): HeLa cells,

## Background Information

CDKN2A generates several transcript variants which differ in their first exons. At least three alternatively-spliced variants encoding distinct proteins were reported. Two of them named p16-INK4 and p14 are sharing 50% identity. The third one -p14(ARF) is entirely unrelated. 10883-1-AP reacts with p16 specifically. P16 plays an essential role in regulating the cell cycle, and mutations in p16 increase the risk of developing various cancers, including melanoma.

## Notable Publications

Author	Pubmed ID	Journal	Application
Shin Hamada	28971839	Am J Physiol Gastrointest Liver Physiol	
Julie Wang	26416809	Circulation	WB
Shengya Tian	31562192	Life Sci Alliance	WB

## Storage

### Storage:

Store at -20°C. Stable for one year after shipment.

### Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)


W: [ptgcn.com](http://ptgcn.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 10883-1-AP (P16-INK4A antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.




WB result of p16-INK4A antibody (10883-1-AP, 1:2,000) with si-Control and si-p16 transfected HEK-293 cells.




Immunohistochemical analysis of paraffin-embedded human cervical cancer tissue slide using 10883-1-AP (P16-INK4A antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded human cervical cancer tissue slide using 10883-1-AP (P16-INK4A antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1x10<sup>6</sup> HeLa cells were intracellularly stained with 0.4 ug Anti-Human P16-INK4A (10883-1-AP)(red), or 0.4 ug Rabbit IgG control Rabbit PolyAb (30000-0-AP) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).



IP result of anti-P16-INK4A (IP:10883-1-AP, 4ug; Detection:10883-1-AP 1:5000) with HEK-293 cells lysate 1560 ug.



Immunofluorescent analysis of (4% PFA) fixed MDCK cells using P16-INK4A antibody (10883-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-phalloidin (red).