

For Research Use Only

NCOA4 Polyclonal antibody

Catalog Number:10968-1-AP

Featured Product

6 Publications



Basic Information

Catalog Number: 10968-1-AP	GenBank Accession Number: BC012736	Purification Method: Antigen affinity purification
Size: 500 ug/ml	GeneID (NCBI): 8031	Recommended Dilutions: IHC 1:50-1:500
Source: Rabbit	UNIPROT ID: Q13772	
Isotype: IgG	Full Name: nuclear receptor coactivator 4	
Immunogen Catalog Number: AG1408	Calculated MW: 70 kDa	

Applications

Tested Applications: IHC, ELISA	Positive Controls: IHC : mouse brain tissue,
Cited Applications: WB, IHC, IF	
Species Specificity: human, mouse	
Cited Species: human, mouse	
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

Nuclear receptor coactivator 4 (NCOA4) also named androgen receptor (AR) coactivator ARA70, RFG and ELE1, is a putative co-activator that specifically enhances the activity of the androgen receptor. In human thyroid carcinomas, the Ret proto-oncogene fuses to ARA70 to form Ret/PTC3 by an intrachromosomal inversion of chromosome 10 in vivo. ARA70a can function as a ligand-enhanced co-activator of PPARγ in adipocytes. However, PPARγ-ARA70 transactivation can be squelched by AR, which suggests cross talk between PPARγ- and AR-mediated response. ARA70a has no intrinsic transcription activation domain or histone acetyltransferase activity, but it interacts with histone acetyltransferase, p/CAF, CBP and p300/CBP-associated factors, and the basal transcription factor TFIIIB. The interaction between ARA70 and AR occurs through the ligand-binding domain. The presence of ARA70 can enhance the androgenic activity of 17-β Estradiol (E2) and antiandrogens toward AR. ARA70 may be involved in prostate carcinogenesis and ovarian cancer and may serve as a key mediator of estrogen-androgen synergism. There are several isoforms of ZGPAT that are produced as a result of alternative splicing events. ARA70a is widely expressed, and its expression is highest in testis and adipose tissues, whereas ARA70b (The shorter variant, results from an internal 985-bp deletion.) is solely expressed in the testis. This antibody is a rabbit polyclonal antibody raised against the N-400aa of human NCOA4 protein.

Notable Publications

Author	Pubmed ID	Journal	Application
Hao Li	34917147	J Oncol	WB,IF
Chun Pan	39366441	Environ Res	WB
Yini Shang	38945330	Int J Biol Macromol	WB,IHC


Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:
T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 10968-1-AP (NCOA4 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 10968-1-AP (NCOA4 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed MCF-7 cells using NCOA4 antibody (10968-1-AP) at dilution of 1:400 and Multi-rAb CoraLite © Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002).