## For Research Use Only

## PROX1 Polyclonal antibody

Catalog Number: 11067-2-AP

Featured Product

43 Publications




## Catalog Number: GenBank Accession Number: **Purification Method: Basic Information** 11067-2-AP BC024201 Antigen affinity purification GenelD (NCBI): Recommended Dilutions: Size: 650 µg/ml 5629 WB 1:2000-1:10000 IF 1:20-1:200 UNIPROT ID: Source: Rabbit Q92786 Full Name: Isotype: IgG prospero homeobox 1 Calculated MW: Immunogen Catalog Number: AG1543 737 aa, 83 kDa **Observed MW:** 90 kDa **Applications Tested Applications:** Positive Controls: FC, IF/ICC, WB, ELISA WB: HepG2 cells, human liver tissue, MCF-7 cells, **Cited Applications:** HuH-7 cells ChIP, FC, IF, IP, WB IF : HepG2 cells, **Species Specificity:** human **Cited Species:** human, rat, mouse, pig **Background Information** PROX1, Prospero homeobox protein 1, belongs to the Prospero homeobox family and contain a Prospero-type homeobox DNA-binding domain. PROX1 plays a fundamental role in early development of CNS and regulates the expression of gene and development of postmitotic, undifferentiated neurons. Also, PROX1 is a required regulator of the development of the lymphatic system. This antibody is raised against part of C-terminal chain of human PROX1. The calculated molecular weight of PROX1 is 83 kDa, but with post-translational modifications, PROX1 is about 90 kDa. PROX1 can be detected double bands with sumo modifications (PMID: 18948102). **Notable Publications** Author Pubmed ID Journal Application LinXi Cheng 34489520 Commun Biol IF Yoshiya Asano 28875528 J Tissue Eng Regen Med Juliana Falero-Perez 30372497 PLoS One **Storage** Storage: Store at -20°C. Stable for one year after shipment. Storage Buffer PBS with 0.02% sodium azide and 50% glycerol pH 7.3. Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 11067-2-AP (PROX1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of HepG2 cells, using PROX1 antibody 11067-2-AP at 1:100 dilution and FITC-labeled donkey anti-rabbit IgG(green). Blue pseudocolor = DAPI (fluorescent DNA dye).



1X10^6 HepG2 cells were stained with 0.2ug PROX1 antibody (11067-2-AP, red) and control antibody (blue). Fixed with 90% MeOH blocked with 3% BSA (30 min). Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) with dilution 1:1000.