## For Research Use Only

# IFT57 Polyclonal antibody

Catalog Number: 11083-1-AP

**Featured Product** 

21 Publications



**Basic Information** 

Catalog Number: GenBank Accession Number: 11083-1-AP BC011899 GeneID (NCBI): Size: 500 μg/ml 55081 **UNIPROT ID:** Source: Rabbit Q9NWB7 Full Name: Isotype:

intraflagellar transport 57 homolog (Chlamydomonas)

Immunogen Catalog Number: AG1566 Calculated MW: 49 kDa

> Observed MW: 49 kDa

**Purification Method:** Antigen affinity purification Recommended Dilutions: WB 1:500-1:1000

> IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate IHC 1:50-1:500 IF 1:50-1:500

**Applications** 

**Tested Applications:** IF/ICC, IHC, IP, WB, ELISA

**Cited Applications:** IF, IP, WB

Species Specificity: human, mouse, rat, Canine

**Cited Species:** 

human, Chicken, rat, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: mouse retina tissue, HEK-293 cells

IP: HEK-293 cells,

IHC: human prostate cancer tissue, human pancreas

tissue

IF: MDCK cells,

# **Background Information**

 $IFT57, also \ named \ as \ DERP8, ESRRBL1, HIPPI \ and \ MHS4R2, belongs \ to \ the \ IFT57 \ family. \ It \ is \ required \ for \ the \ IFT57 \ family.$ formation of cilia. IFT57 plays an indirect role in sonic hedgehog signaling, cilia being required for all activity of the hedgehog pathway. It is a component of IFT complex B which composed of IFT88, IFT57, TRAF3IP1, IFT52, IFT27, HSPB11 and IFT20. In contrast to BBS proteins, IFT57 and IFT88 is found in almost every cilium. (PMID:22072986)

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Malavika Raman	26389662	Nat Cell Biol	WB
Ying Hsu	29049287	PLoS Genet	WB
Laura Vuolo	30320547	Elife	WB,IF

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


### **Selected Validation Data**




mouse retina tissue were subjected to SDS PAGE followed by western blot with 11083-1-AP (IFT57 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




WB result of IFT57 antibody (11083-1-AP; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-IFT57 transfected HEK-293 cells.



Immunofluorescent analysis of MDCK cells using 11083-1-AP (IFT57 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



IP result of anti-IFT57 (IP:11083-1-AP, 5ug; Detection:11083-1-AP 1:800) with HEK-293 cells lysate 2000ug.



Immunohistochemical analysis of paraffinembedded human prostate cancer using 11083-1-AP (IFT57 antibody) at dilution of 1:50 (under 10x lens)