

For Research Use Only

NEK9 Polyclonal antibody

Catalog Number: 11192-1-AP **2 Publications**



Basic Information

Catalog Number: 11192-1-AP	GenBank Accession Number: BC009336	Purification Method: Antigen affinity purification
Size: 750 µg/ml	GeneID (NCBI): 91754	Recommended Dilutions: WB 1:1000-1:9000
Source: Rabbit	UNIPROT ID: Q8TD19	IP 0.5-4.0 µg for 1.0-3.0 mg of total protein lysate
Isotype: IgG	Full Name: NIMA (never in mitosis gene a)-related kinase 9	IHC 1:50-1:500
Immunogen Catalog Number: AG1678	Calculated MW: 107 kDa	IF/ICC 1:200-1:800
	Observed MW: 120 kDa	

Applications

Tested Applications: WB, IHC, IF/ICC, FC (Intra), IP, ELISA	Positive Controls: WB : HeLa cells, human placenta tissue, HepG2 cells
Cited Applications: WB, IF	IP : HeLa cells,
Species Specificity: human, mouse, rat	IHC : human lymphoma tissue,
Cited Species: human	IF/ICC : HepG2 cells,
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

NEK9 is also named as KIAA1995, NEK8, NERCC and belongs to the NEK Ser/Thr protein kinase family. It is originally identified as α β -casein kinase that associates with a putative substrate Bicc2. It is also independently identified as a Nek6- and Ran GTPase-binding protein under a different name, Nerc1(PMID:14660563). Nek9, together with the highly similar (80% identical) Nek6 and Nek7, form a signaling module that is activated during mitosis and involved in the regulation of the mitotic spindle. The endogenous Nek9 (with predicted molecular mass of 120 kDa) has an apparent molecular mass of 600 kDa, also compatible with a tetramer(PMID:21454704).

Notable Publications

Author	Pubmed ID	Journal	Application
Andrea M Brum	30283877	JBMR Plus	IF
Lei Chen	37835513	Cancers (Basel)	WB

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Immunohistochemical analysis of paraffin-embedded human lymphoma tissue slide using 11192-1-AP (NEK9 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Various lysates were subjected to SDS PAGE followed by western blot with 11192-1-AP (NEK9 antibody) at dilution of 1:4500 incubated at room temperature for 1.5 hours.




IP result of anti-NEK9 (IP:11192-1-AP, 4ug; Detection:11192-1-AP 1:1000) with HeLa cells lysate 1880 ug.



Immunofluorescent analysis of HepG2 cells, using NEK9 antibody 11192-1-AP at 1:50 dilution and FITC-labeled donkey anti-rabbit IgG (green). Blue pseudocolor = DAPI (fluorescent DNA dye).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using NEK9 antibody (11192-1-AP) at dilution of 1:400 and CoraLite@488-Conjugated AffiniPure Goat Anti-Rabbit IgG (H+L).



1×10^6 HeLa cells were intracellularly stained with 0.4 ug Anti-Human NEK9 (11192-1-AP) and CoraLite@488-Conjugated AffiniPure Goat Anti-Rabbit IgG (H+L) at dilution 1:1000 (red), or 0.4 ug x. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).