

For Research Use Only

# HMGN1 Polyclonal antibody

Catalog Number: 11695-1-AP

8 Publications



## Basic Information

<b>Catalog Number:</b> 11695-1-AP	<b>GenBank Accession Number:</b> BC023984	<b>Purification Method:</b> Antigen affinity purification
<b>Size:</b> 1000 ug/ml	<b>GeneID (NCBI):</b> 3150	<b>Recommended Dilutions:</b> WB 1:500-1:2000 IHC 1:50-1:500 IF/ICC 1:200-1:800
<b>Source:</b> Rabbit	<b>UNIPROT ID:</b> P05114	
<b>Isotype:</b> IgG	<b>Full Name:</b> high-mobility group nucleosome binding domain 1	
<b>Immunogen Catalog Number:</b> AG2315	<b>Calculated MW:</b> 100 aa, 11 kDa	
	<b>Observed MW:</b> 17 kDa	

## Applications

<b>Tested Applications:</b> WB, IHC, IF/ICC, ELISA	<b>Positive Controls:</b> WB : HepG2 cells, HeLa cells
<b>Cited Applications:</b> WB, IHC	<b>IHC :</b> human kidney tissue, mouse kidney tissue
<b>Species Specificity:</b> human, mouse, rat	<b>IF/ICC :</b> HepG2 cells,
<b>Cited Species:</b> human, mouse	

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

## Background Information

HMGN1 (high-mobility group nucleosome-binding protein 1) is a member of the HMG superfamily of nonhistone chromatin-binding proteins, which which regulate chromatin structure and gene transcription. It binds to the inner side of the nucleosomal DNA thus altering the interaction between the DNA and the histone octamer. It may participate in the process which maintains transcribable genes in an unique chromatin conformation, and inhibit the phosphorylation of nucleosomal histones H3 and H2A by RPS6KA5/MSK1 and RPS6KA3/RSK2 expressed abundantly

## Notable Publications

Author	Pubmed ID	Journal	Application
Feng Wei	25205103	Cancer Res	WB
Zhili Liu	31281528	Theranostics	WB
Dan Zhang	30031066	J Proteomics	WB

## Storage

**Storage:**  
Store at -20°C. Stable for one year after shipment.  
**Storage Buffer:**  
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.  
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:

T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**


## Selected Validation Data




HepG2 cells were subjected to SDS PAGE followed by western blot with 11695-1-AP (HMGN1 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human kidney tissue slide using 11695-1-AP (HMGN1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded human kidney tissue slide using 11695-1-AP (HMGN1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse kidney tissue slide using 11695-1-AP (HMGN1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse kidney tissue slide using 11695-1-AP (HMGN1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using HMGN1 antibody (11695-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-phalloidin (red).