## For Research Use Only

# ASGR1 Polyclonal antibody

Catalog Number: 11739-1-AP

Featured Product

16 Publications

BC032130

GenBank Accession Number:



**Basic Information** 

Catalog Number: 11739-1-AP

Size: GeneID (NCBI):

650 μg/ml 43

Source: UNIPROT ID:
Rabbit P07306

Isotype: Full Name:

gG asialoglycoprotein receptor 1

Immunogen Catalog Number:Calculated MW:AG2310291 aa, 33 kDa

Observed MW: 40 kDa Purification Method:

Antigen affinity purification

Recommended Dilutions: WB 1:5000-1:50000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:1000-1:4000 IF 1:50-1:500

**Applications** 

**Tested Applications:** 

FC, IF-P, IHC, IP, WB, ELISA

Cited Applications: CoIP, FC, IF, IHC, WB Species Specificity: human, mouse, rat Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 **Positive Controls:** 

WB: HepG2 cells, human liver tissue, mouse liver tissue, rat liver tissue

IP: mouse liver tissue,

IHC: rat liver tissue, human liver cancer tissue

IF: human liver cancer tissue, rat liver tissue

# Background Information

Asialoglycoprotein receptor (ASGPR), also known as the hepatic galactose/N-acetylglucosamine (GlcNAc) receptor or Ashwell receptor, is a C-type lectin expressed exclusively in hepatic parenchymal cells. ASGPR consists of two subunits, a major subunit (ASGR1, HL-1) and a minor subunit (ASGR2, HL-2), and specifically recognizes terminal  $\beta$ -linked galactose or GlcNAc on circulating glycoproteins or cells. This receptor plays a critical role in serum glycoprotein homeostasis by mediating the endocytosis and lysosomal degradation of glycoproteins that contain terminal galactose or GlcNAc residues. ASGPR may facilitate hepatic infection by multiple viruses including hepatitis B, and is also a target for liver-specific drug delivery.

#### **Notable Publications**

| Author           | Pubmed ID | Journal                | Application |
|------------------|-----------|------------------------|-------------|
| Juergen Scharner | 30965276  | Mol Ther Nucleic Acids | WB,IHC      |
| Wai Kit Ma       | 34921016  | Cancer Res             | WB          |
| Youngsoo Kim     | 31303442  | Mol Ther               | IHC,IF      |

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926 E: Proteintech-CN@ptglab.com

W: ptgcn.co

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 11739-1-AP (ASGR1 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded rat liver tissue slide using 11739-1-AP (ASGR1 antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded rat liver tissue slide using 11739-1-AP (ASGR1 antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




IP result of anti-ASGR1 (IP:11739-1-AP, 4ug; Detection:11739-1-AP 1:2000) with mouse liver tissue lysate 4000 ug.



Immunofluorescent analysis of (4% PFA) fixed human liver cancer tissue using ASGR1 antibody (11739-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed human liver cancer tissue using ASGR1 antibody (11739-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



1X10^6 HepG2 cells were intracellularly stained with 0.4 ug Anti-Human ASGR1 (11739-1-AP) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).