For Research Use Only

RAMP3 Polyclonal antibody

Catalog Number: 11861-1-AP

1 Publications



Basic Information

Catalog Number: GenBank Accession Number: 11861-1-AP BC022304 GeneID (NCBI): Size: 200 μg/ml 10268 **UNIPROT ID:** Source: Rabbit 060896 Full Name: Isotype:

receptor (G protein-coupled) activity

Immunogen Catalog Number: AG2438 Calculated MW: 16.5 kDa Observed MW:

45-50 kDa

modifying protein 3

Applications

Tested Applications: IHC, WB, ELISA **Cited Applications:**

Species Specificity: human, mouse Cited Species:

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: A549 cells, human kidney tissue, Jurkat cells, mouse pancreas tissue, human placenta tissue, mouse brain tissue, mouse heart tissue, mouse skeletal muscle tissue, human heart tissue, mouse lung tissue, MCF-7 cells

Purification Method:

WB 1:500-1:1000 IHC 1:20-1:200

Antigen affinity purification

Recommended Dilutions:

IHC: human breast cancer tissue,

Background Information

 $The \ receptor \ activity-modifying \ proteins \ (RAMPs) \ and \ the \ calciton in \ receptor-like \ receptor \ (CRLR) \ are \ both \ required$ to generate adrenomedullin (AM) and calcitonin gene-related peptide (CGRP) receptors. During the process, RAMP3 transports the glycosylated 58-kD CGRPR to the cell surface. The formation of CRLR-RAMP heterodimer is essential for the proper cell surface targeting and pharmacological characteristics of both CGRP and AM receptors.

Notable Publications

Author	Pubmed ID	Journal	Application
Zao-Ling Liu	38946034	J Chin Med Assoc	IHC

Storage


Storage:

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 11861-1-AP (RAMP3 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human breast cancer using 11861-1-AP (RAMP3 antibody) at dilution of 1:50 (under 10x lens).