

For Research Use Only

# XIRP2 Polyclonal antibody

Catalog Number: 11896-1-AP

3 Publications



## Basic Information

<b>Catalog Number:</b> 11896-1-AP	<b>GenBank Accession Number:</b> BC022888	<b>Purification Method:</b> Antigen affinity purification
<b>Size:</b> 240 µg/ml	<b>GeneID (NCBI):</b> 129446	<b>Recommended Dilutions:</b> WB 1:500-1:1000 IHC 1:20-1:200 IF-P 1:50-1:500
<b>Source:</b> Rabbit	<b>UNIPROT ID:</b> A4UGR9	
<b>Isotype:</b> IgG	<b>Full Name:</b> xin actin-binding repeat containing 2	
<b>Immunogen Catalog Number:</b> AG2515	<b>Calculated MW:</b> 379 kDa	
	<b>Observed MW:</b> 260-300 kDa	

## Applications

<b>Tested Applications:</b> WB, IHC, IF-P, ELISA	<b>Positive Controls:</b> WB : COLO 320 cells, Caco-2 cells, human heart tissue IHC : human skeletal muscle tissue, human heart tissue IF-P : mouse heart tissue,
<b>Cited Applications:</b> WB, IHC, IF	
<b>Species Specificity:</b> human, mouse	
<b>Cited Species:</b> human, mouse	

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

## Background Information

### Notable Publications

Author	Pubmed ID	Journal	Application
Emily Farrell	30410445	Front Physiol	WB,IF
Dahuan Li	39194571	Biology (Basel)	WB,IF
Elizabeth L Wagner	37294664	Elife	WB,IHC

## Storage

**Storage:**  
Store at -20°C. Stable for one year after shipment.  
**Storage Buffer:**  
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.  
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:

T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)


W: [ptgcn.com](http://ptgcn.com)

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**

## Selected Validation Data




COLO 320 cells were subjected to SDS PAGE followed by western blot with 11896-1-AP (XIRP2 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human skeletal muscle using 11896-1-AP (XIRP2 antibody) at dilution of 1:50 (under 10x lens).



Immunohistochemical analysis of paraffin-embedded human skeletal muscle using 11896-1-AP (XIRP2 antibody) at dilution of 1:50 (under 40x lens).



Immunofluorescent analysis of (4% PFA) fixed mouse heart tissue using XIRP2 antibody (11896-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L).