

For Research Use Only

HELLS Polyclonal antibody

Catalog Number: 11955-1-AP **11 Publications**



Basic Information

Catalog Number: 11955-1-AP	GenBank Accession Number: BC031004	Purification Method: Antigen affinity purification
Size: 750 ug/ml	GeneID (NCBI): 3070	Recommended Dilutions: WB 1:1000-1:5000
Source: Rabbit	UNIPROT ID: Q9NRZ9	IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate
Isotype: IgG	Full Name: helicase, lymphoid-specific	IHC 1:500-1:2000
Immunogen Catalog Number: AG2558	Calculated MW: 348 aa, 40 kDa, 97 kDa	IF/ICC 1:200-1:800
	Observed MW: 97 kDa	

Applications

Tested Applications: WB, IHC, IF/ICC, FC (Intra), IP, ELISA	Positive Controls: WB : K-562 cells, DU 145 cells, mouse testis tissue, mouse thymus tissue, PC-3 cells
Cited Applications: WB, IF	IP : mouse testis tissue,
Species Specificity: human, mouse, rat	IHC : mouse thymus tissue, mouse testis tissue, rat testis tissue, rat thymus tissue
Cited Species: human, mouse	IF/ICC : HepG2 cells,

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

Lymphoid-specific helicase(HELLS) is a member of the SNF2 family of chromatin remodeling proteins that is required for development and survival. It's required for de novo or maintenance DNA methylation, which mediated the silencing of the imprinted CDKN1C gene. Also it involved in formation and organization of heterochromatin that regulate transcription and mitosis.

Notable Publications

Author	Pubmed ID	Journal	Application
Xuan Yang	31066149	Cancer Sci	WB
Feng-Jiao Wang	34315468	BMC Med Genomics	WB
Haoyu Wu	27466202	Hum Mol Genet	


Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:
T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




K-562 cells were subjected to SDS PAGE followed by western blot with 11955-1-AP (HELLS antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.




IP result of anti-HELLS (IP:11955-1-AP, 4ug; Detection:11955-1-AP 1:1000) with mouse testis tissue lysate 4000ug.




Immunohistochemical analysis of paraffin-embedded mouse thymus tissue slide using 11955-1-AP (HELLS antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded mouse thymus tissue slide using 11955-1-AP (HELLS antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded mouse testis tissue slide using 11955-1-AP (HELLS antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded mouse testis tissue slide using 11955-1-AP (HELLS antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).






Immunohistochemical analysis of paraffin-embedded rat testis tissue slide using 11955-1-AP (HELLS antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded rat testis tissue slide using 11955-1-AP (HELLS antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded rat thymus tissue slide using 11955-1-AP (HELLS antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded rat thymus tissue slide using 11955-1-AP (HELLS antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).

Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using HELLs antibody (11955-1-AP) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red).

1×10^6 HepG2 cells were intracellularly stained with 0.4 ug HELLs Polyclonal antibody (11955-1-AP) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2)(red), or 0.4 ug Rabbit IgG control Rabbit PolyAb (30000-0-AP) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).