For Research Use Only

SUB1 Polyclonal antibody

Catalog Number: 11956-1-AP

Featured Product



Basic Information

Catalog Number: GenBank Accession Number: 11956-1-AP BC022339 GeneID (NCBI): Size: 1000 µg/ml 10923 Source: **UNIPROT ID:** Rabbit P53999 Isotype: Full Name:

SUB1 homolog (S. cerevisiae)

Calculated MW: Immunogen Catalog Number:

AG2559 14 kDa

> Observed MW: 18-23 kDa

Applications

Tested Applications: WB, IHC, IF/ICC, ELISA Species Specificity:

human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Recommended Dilutions: WB 1:500-1:2000 IHC 1:50-1:500 IF/ICC 1:50-1:500

Purification Method:

Antigen affinity purification

Positive Controls:

WB: rat liver tissue, HeLa cells, C2C12 cells

IHC: human gliomas tissue,

IF/ICC: HeLa cells,

Background Information

SUB1 gene encodes activated RNA polymerase II transcriptional coactivator p15, also termed as positive cofactor 4 (PC4) or p14. SUB1 is a sort of general coactivator that functions in assembling and stablization of transcription complex. It has DNA binding activity and is localized in the nucleus, mulitple modification sites of phosphorylation $or a cetylation\ may\ turn\ it\ appear\ slightly\ shifted\ molecular\ weight.\ Catalog \#11956-1-AP\ is\ a\ rabbit\ polyclonal\ molecular\ weight.$ antibody raised against full-length human SUB1. This antibody recognize 14-16 kDa (P14 or P15), 20 kDa(phospho form) and 30 kDa (Dimer form).

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Immunohistochemical analysis of paraffinembedded human gliomas tissue slide using 11956-1-AP (SUB1 antibody) at dilution of 1:100 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



rat liver tissue were subjected to SDS PAGE followed by western blot with 11956-1-AP (SUB1 Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (10% Formaldehyde) fixed HeLa cells using 11956-1-AP (SUB1 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



WB result of SUB1 antibody (11956-1-AP; 1:3000; incubated at room temperature for 1.5 hours) with sh-Control and sh-SUB1 transfected HeLa cells.