

For Research Use Only

SERPING1/C1 Inactivator Polyclonal antibody

Catalog Number: 12259-1-AP

9 Publications



Basic Information

Catalog Number:

12259-1-AP

Size:

500 ug/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG2819

GenBank Accession Number:

BC011171

GeneID (NCBI):

710

UNIPROT ID:

P05155

Full Name:

serpin peptidase inhibitor, clade G (C1 inhibitor), member 1

Calculated MW:

500 aa, 55 kDa

Observed MW:

100 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:1000-1:4000

IHC 1:200-1:800

IF/ICC 1:200-1:800

Applications

Tested Applications:

WB, IHC, IF/ICC, ELISA

Cited Applications:

WB, IHC, IF

Species Specificity:

human, mouse

Cited Species:

human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: human plasma, HeLa cells, human liver tissue, L02 cells, mouse liver tissue, HepG2 cells, human urine tissue

IHC: human tonsillitis tissue, human normal colon

IF/ICC: HepG2 cells,

Background Information

SERPING1, also known as C1 Inhibitor (C1INH), is a member of the Serine proteinase inhibitor family. Its main function is the inhibition of the complement system to prevent spontaneous activation. It inhibits activated C1r and C1s of the first complement component and thus regulates complement activation. Deficiency of this protein is associated with hereditary angioneurotic oedema (HANE). C1INH is the most heavily glycosylated plasma protein. Of its 95-105 kDa apparent molecular mass, the protein moiety of 500 aa accounts for only 55 kDa. The native 105 kDa form of C1INH could be cleaved into 60-65 kDa fragments.

Notable Publications

Author	Pubmed ID	Journal	Application
Daria Zamolodchikov	26613657	J Thromb Haemost	WB
Koji Abe	31771070	J Alzheimers Dis	IHC
Dorit Farfara	30882931	Glia	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 12259-1-AP (SERPING1/C1 Inactivator antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 12259-1-AP (SERPING1/C1 Inactivator antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 12259-1-AP (SERPING1/C1 Inactivator antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using SERPING1/C1 Inactivator antibody (12259-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).



Immunohistochemical analysis of paraffin-embedded human normal colon slide using 12259-1-AP (SERPING1/C1 Inactivator antibody) at dilution of 1:400 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human normal colon slide using 12259-1-AP (SERPING1/C1 Inactivator antibody) at dilution of 1:400 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).