For Research Use Only

Sortilin Polyclonal antibody

Catalog Number: 12369-1-AP

Featured Product

14 Publications



Basic Information

Catalog Number: 12369-1-AP

300 μg/ml

Rabbit Isotype:

Immunogen Catalog Number:

AG2997

Size:

Source:

GenBank Accession Number:

BC023542 GeneID (NCBI): 6272

ENSEMBL Gene ID:

ENSG00000134243 UNIPROT ID: Q99523 Full Name: sortilin 1

Calculated MW: 831 aa, 92 kDa

Observed MW: 100-110 kDa

Applications

Tested Applications:

WB, IHC, IF-P, FC (Intra), IP, ELISA

Cited Applications: WB, IHC, IF, IP, CoIP Species Specificity: human, mouse, rat Cited Species:

human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: mouse brain tissue, human brain tissue, rat brain

Purification Method:

WB 1:2000-1:16000

protein lysate

IHC 1:500-1:2000

IF-P 1:200-1:800

Antigen affinity purification

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

tissue

IP: rat brain tissue,

IHC: mouse brain tissue, human brain tissue, rat brain

tissue

IF-P: mouse brain tissue,

Background Information

Sortilin 1 (SORT1), is a trans-Golgi network (TGN) transmembrane protein and is a member of the Vps10p domain receptor family. SORT1 binds a number of unrelated ligands that participate in a wide range of cellular processes. It functions as a sorting receptor in the Golgi compartment and as a clearance receptor on the cell surface.

Notable Publications

Author	Pubmed ID	Journal	Application
Hiroyuki Obata	36185767	Exp Ther Med	IF
Keiji Uchiyama	28665987	PLoS Pathog	WB,IF
Noah R Johnson	29203673	Proc Natl Acad Sci U S A	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 12369-1-AP (Sortilin antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 12369-1-AP (Sortilin antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 12369-1-AP (Sortilin antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




IP result of anti-Sortilin (IP:12369-1-AP, 4ug; Detection:12369-1-AP 1:500) with rat brain tissue lysate 4000ug.



Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using Sortilin antibody (12369-1-AP) at dilution of 1:400 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using Sortilin antibody (12369-1-AP) at dilution of 1:400 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L).



1X10^6 Neuro-2a cells were intracellularly stained with 0.4 ug Anti-Human Sortilin (12369-1-AP) and Coralite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Isotype Control. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).