

For Research Use Only

TSKU Polyclonal antibody

Catalog Number: 12370-1-AP



Basic Information

| | | |
|--|--|--|
| Catalog Number: 12370-1-AP | GenBank Accession Number: BC020975 | Purification Method: Antigen affinity purification |
| Size: 550 µg/ml | GeneID (NCBI): 25987 | Recommended Dilutions: WB 1:200-1:1000 IHC 1:50-1:500 |
| Source: Rabbit | UNIPROT ID: Q8WUA8 | |
| Isotype: IgG | Full Name: tsukushin | |
| Immunogen Catalog Number: AG3054 | Calculated MW: 353 aa, 38 kDa | |
| | Observed MW: 45-50 kDa | |

Applications

Tested Applications:

IHC, WB, ELISA

Species Specificity:

human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : mouse liver tissue, HepG2 cells, mouse colon tissue, mouse pancreas tissue

IHC : human colon cancer tissue, human breast cancer tissue

Background Information

TSK, also known as Tsukushi, is an atypical member of the small leucine-rich proteoglycan (SLRP) family of extracellular matrix proteins (PMID: 16884925). The human TSK cDNA encodes a 16 amino acid (aa) signal sequence and a 337 aa secreted protein that contains nine LRR, which has a calculated molecular mass of 38kDa and an apparent molecular mass of 40-50kDa cause of the N-glycosylation. In chick embryos, C-TSK is a unique BMP inhibitor that forms a ternary complex with BMP and chordin and is involved in the regulation of primitive streak and node formation. C-TSK migrates as a band of 47 kDa in SDS-PAGE and N-glycosidase F treatment reduces the size to 41 kDa (PMID: 15363410).

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



mouse liver tissue were subjected to SDS PAGE followed by western blot with 12370-1-AP (TSKU Antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human colon cancer tissue slide using 12370-1-AP (TSKU Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human colon cancer tissue slide using 12370-1-AP (TSKU Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).