For Research Use Only

RHPN2 Polyclonal antibody

Catalog Number: 12671-1-AP

Featured Product

3 Publications



Basic Information

Catalog Number: GenBank Accession Number: 12671-1-AP BC036447 GeneID (NCBI): Size: 500 ug/ml 85415 **UNIPROT ID:** Source: Rabbit Q8IUC4 Full Name: Isotype:

Antigen affinity purification Recommended Dilutions: WB 1:500-1:1000 IHC 1:50-1:500 IF/ICC 1:200-1:800

Purification Method:

rhophilin, Rho GTPase binding protein

Immunogen Catalog Number:

Calculated MW:

686 aa. 77 kDa Observed MW: 77 kDa

Applications

Tested Applications: WB, IHC, IF/ICC, ELISA Cited Applications: WB, IF

Species Specificity: human, mouse, rat **Cited Species:** human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: COLO 320 cells, HepG2 cells

IHC: human colon tissue, human stomach tissue

IF/ICC: HepG2 cells,

Background Information

RHPN2 binds specifically to GTP-Rho. It may function in a Rho pathway to limit stress fiber formation and/or increase the turnover of F-actin structures in the absence of high levels of RhoA activity. The MW of this protein is 77 kDa, and this antibody specially recognises the 77 kDa protein.

Notable Publications

Author	Pubmed ID	Journal	Application
Junning Yang	27460990	Sci Signal	
Jian Zhang	36536496	J Biochem Mol Toxicol	WB
Limei Yan	34382920	Cell Cycle	WB,IF


Storage

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Immunohistochemical analysis of paraffinembedded human colon tissue slide using 12671-1-AP (RHPN2 antibody) at dilution of 1:200 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human colon tissue slide using 12671-1-AP (RHPN2 antibody) at dilution of 1:200 (under 40x lens).



COLO 320 cells were subjected to SDS PAGE followed by western blot with 12671-1-AP (RHPN2 Antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using RHPN2 antibody (12671-1-AP) at dilution of 1:400 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002), CL594-phalloidin (red).