

For Research Use Only

DTL Polyclonal antibody

Catalog Number: 12896-1-AP

2 Publications



Basic Information

Catalog Number:

12896-1-AP

Size:

200 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG3589

GenBank Accession Number:

BC033540

GeneID (NCBI):

51514

UNIPROT ID:

Q9NZJ0

Full Name:

denticleless homolog (Drosophila)

Calculated MW:

730 aa, 79 kDa

Observed MW:

79 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:200-1:1000

IP 0.5-4.0 µg for 1.0-3.0 mg of total protein lysate

IF/ICC 1:50-1:500

Applications

Tested Applications:

WB, IF/ICC, IP, ELISA

Cited Applications:

WB, IF

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse

Positive Controls:

WB : HeLa cells, L02 cells, MCF-7 cells

IP : mouse testis tissue,

IF/ICC : HeLa cells,

Background Information

CDT2, also named as L2DTL or RAMP, is one of the substrate receptors of the Cullin Ring Ubiquitin Ligase 4 that targets for ubiquitin mediated degradation a number of substrates, such as CDT1, p21 and CHK1, involved in the regulation of cell cycle and survival. CDT2 overexpression was reported in breast (PMID: 18542055), gastric (PMID: 19672268) and ovarian carcinomas (PMID: 23995842) and rhabdomyosarcomas (PMID: 19235922) and associated with the aggressiveness of hepatocellular carcinomas (PMID: 17106265).

Notable Publications

Author	Pubmed ID	Journal	Application
Chunxia Zhang	35108058	Sci Adv	IF
Chengli Yu	36843329	Proteomics	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



HeLa cells were subjected to SDS PAGE followed by western blot with 12896-1-AP (DTL antibody) at dilution of 1:200 incubated at room temperature for 1.5 hours.



IP result of anti-DTL (IP:12896-1-AP, 3ug; Detection:12896-1-AP 1:500) with mouse testis tissue lysate 8000ug.



Immunofluorescent analysis of (10% Formaldehyde) fixed HeLa cells using 12896-1-AP (DTL antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).