

For Research Use Only

IL-17A Polyclonal antibody

Catalog Number: 13082-1-AP **45 Publications**



Basic Information

Catalog Number: 13082-1-AP	GenBank Accession Number: BC067505	Purification Method: Antigen affinity purification
Size: 650 µg/ml	GeneID (NCBI): 3605	Recommended Dilutions: IHC 1:50-1:500
Source: Rabbit	ENSEMBL Gene ID: ENSG00000112115	
Isotype: IgG	UNIPROT ID: Q16552	
Immunogen Catalog Number: AG3733	Full Name: interleukin 17A	
	Calculated MW: 155 aa, 18 kDa	

Applications

Tested Applications:
IHC, ELISA

Cited Applications:
WB, IF, FC, IHC, Cell treatment

Species Specificity:
human

Cited Species:
human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

IHC: human colon cancer tissue, human cervical cancer tissue, human stomach cancer tissue, human colon tissue

Background Information

IL17A, also named as IL-17, is a proinflammatory cytokine. IL-17, synthesized only by memory T cells and natural killer cells, has pleiotropic effects, mainly in the recruitment and activation of neutrophils. This cytokine regulates the activities of NF-kappaB and mitogen-activated protein kinases. This cytokine can stimulate the expression of IL6 and cyclooxygenase-2 (PTGS2/COX-2), as well as enhance the production of nitric oxide (NO). High levels of this cytokine are associated with several chronic inflammatory diseases including rheumatoid arthritis, psoriasis and multiple sclerosis. The IL-17 receptor is a type I transmembrane protein, that is widely expressed on epithelial cells, fibroblasts, B and T cells, and monocytic cells. In psoriatic skin lesions, both Th17 cells and their downstream effector molecules, e.g. IL-17 and IL-22, are highly increased. This antibody got 32-35 kDa band in western blot, maybe due to homodimer formation and differential glycosylations.

Notable Publications

Author	Pubmed ID	Journal	Application
Yang Liu	36175851	BMC Mol Cell Biol	IF
Yong Xu	34630661	Exp Ther Med	WB
Ying Jiang	34655920	Mol Immunol	WB


Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:
T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Immunohistochemical analysis of paraffin-embedded human colon cancer tissue slide using 13082-1-AP (IL-17A antibody at dilution of 1:200) (under 10x lens).



Immunohistochemical analysis of paraffin-embedded human colon cancer tissue slide using 13082-1-AP (IL-17A antibody at dilution of 1:200) (under 40x lens).