For Research Use Only

CA5B Polyclonal antibody

Catalog Number: 13342-1-AP

1 Publications



Purification Method:

WB 1:1000-1:6000

IHC 1:300-1:1200

Antigen affinity purification Recommended Dilutions:

Basic Information

Catalog Number: 13342-1-AP

Size: 400 µg/ml Source: Rabbit Isotype:

IgG
Immunogen Catalog Number:

AG4166

Observed MW: 32 kDa

Applications

Tested Applications: IHC, WB, ELISA

Cited Applications:

MR

Species Specificity: human, mouse, rat Cited Species: human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

CA5B(Carbonic anhydrase 5B, mitochondrial) is also named as CA-VB and belongs to the alpha-carbonic anhydrase family.CA5B encodes a 317-amino acid protein that has 64% sequence similarity with CA5 (114761), a mitochondrial CA, and contains a 33-amino-acid residue signal sequence. Western blot analysis revealed a 36-kD protein precursor and a 32-kD mature protein(PMID:10409679).

GenBank Accession Number:

carbonic anhydrase VB, mitochondrial

Positive Controls:

WB: mouse testis tissue,

IHC: human kidney tissue,

BC028142

11238

GeneID (NCBI):

UNIPROT ID: Q9Y2D0

Full Name:

Calculated MW:

317 aa, 36 kDa

Notable Publications

AuthorPubmed IDJournalApplicationOliva Palander33872598J Biol ChemWB

Storage

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 13342-1-AP (CA5B antibody) at dilution of 1:600 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 13342-1-AP (CA5B antibody) at dilution of 1:600 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



mouse testis tissue were subjected to SDS PAGE followed by western blot with 13342-1-AP (CA5B antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.