For Research Use Only

## IL-10RA Polyclonal antibody

Catalog Number:13356-1-AP 1 Publications



Basic Information	Catalog Number: 13356-1-AP	GenBank Accession Number: BC028082	Purification Method:	
	Size:	GenelD (NCBI):	Antigen affinity purification Recommended Dilutions:	
	1000 µg/ml	3587	WB 1:500-1:2000	
	Source: Rabbit	UNIPROT ID: Q13651		
	lsotype: IgG	Full Name: interleukin 10 receptor, alpha		
	Immunogen Catalog Number: AG4156	Calculated MW: 578 aa, 63 kDa		
		Observed MW: 70-72 kDa, 90-95 kDa		
Applications	Tested Applications:	Positive (	tive Controls: HeLa cells, human heart tissue, human placenta e, human liver tissue, K-562 cells, mouse brain e, rat brain tissue, rat liver tissue	
	WB, ELISA Cited Applications:			
	IF	•		
	Species Specificity: human, mouse, rat			
	Cited Species: mouse			
		Interleukin-10 (IL-10) is an anti-inflammatory cytokine that plays a critical role in maintaining immune system balance. IL-10 signals through a receptor complex consisting of two molecules of the IL-10R alpha-chain (IL-10RA) and two molecules of the accessory IL-10R beta-chain (IL-10RB) (PMID: 22236434). IL-10RA and IL-10RB are member of the type II cytokine receptor family. IL-10RA, also known as IL-10R1 or CD210, is expressed on most hematopoietic cells at a basal level but is upregulated by various cells upon activation (PMID: 24507158). IL-10RA serves as the ligand-binding subunit of the receptor complex. Upon binding of IL-10 to the IL-10R complex, the kinases JAK1 and TYK2 are activated and catalyze the phosphorylation of STAT3 (PMID: 11244051; 10433356).		
Background Information	and two molecules of the accesso of the type II cytokine receptor fai hematopoietic cells at a basal lev serves as the ligand-binding subu kinases JAK1 and TYK2 are activa	eceptor complex consisting of two mo ry IL-10R beta-chain (IL-10RB) (PMID: : mily. IL-10RA, also known as IL-10R1 rel but is upregulated by various cells nit of the receptor complex. Upon bin ted and catalyze the phosphorylation	lecules of the IL-10R alpha-chain (IL-10RA) 22236434). IL-10RA and IL-10RB are membe or CD210, is expressed on most upon activation (PMID: 24507158). IL-10RA ding of IL-10 to the IL-10R complex, the of IL-10RA at specific intracellular tyrosine	
Background Information	and two molecules of the accesso of the type II cytokine receptor fa hematopoietic cells at a basal lev serves as the ligand-binding subu kinases JAK1 and TYK2 are activa residues, leading to the recruitme	eceptor complex consisting of two mo ry IL-10R beta-chain (IL-10RB) (PMID: : mily. IL-10RA, also known as IL-10R1 rel but is upregulated by various cells nit of the receptor complex. Upon bin ted and catalyze the phosphorylation	lecules of the IL-10R alpha-chain (IL-10RA) 22236434). IL-10RA and IL-10RB are member or CD210, is expressed on most upon activation (PMID: 24507158). IL-10RA ding of IL-10 to the IL-10R complex, the of IL-10RA at specific intracellular tyrosine	

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 13356-1-AP (IL-10RA antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.