For Research Use Only

CD86 Polyclonal antibody

Catalog Number:13395-1-AP

140 Publications




Basic Information	Catalog Number:GenBank Accession Number:13395-1-APBC040261		Number:	Purification Method: Antigen affinity purification	
	Size:	GeneID (NCBI):		Recommended Dilutions:	
	500 µg/ml 942			WB 1:1000-1:4000 IF 1:50-1:500	
	Source: Rabbit	ENSEMBL Gene ID: ENSG00000114013		1 1.90 1.900	
	lsotype: IgG	UNIPROT ID: P42081			
	Immunogen Catalog Number:Full Name:AG4063CD86 molecule				
	Calculated MW: 329 aa, 38 kDa				
	Observed MW: 60-80 kDa				
Applications	Tested Applications:		Positive Controls:		
	FC, IF/ICC, WB, ELISA Cited Applications:		WB : Daudi cells, Raji cells, Ramos cells, RAW 264.7 cells		
	WB, IF, FC, IHC, Cell treatment		IF : raji cells,		
	Species Specificity: human, mouse		in Tuji cetto,		
	Cited Species: human, rat, mouse				
Background Information	CD86 (also known as B7.2) is a costimulatory molecule belonging to the immunoglobulin superfamily. Primarily expressed on antigen-presenting cells (APCs), including B cells, dendritic cells, and macrophages, CD86 is the ligand for two proteins at the cell surface of T cells, CD28 antigen and cytotoxic T-lymphocyte-associated protein 4. Binding of CD86 with CD28 antigen is a costimulatory signal for activation of the T-cell. Binding of CD86 with cytotoxic T-lymphocyte-associated protein 4 negatively regulates T-cell activation and diminishes the immune response. This antibody detects a band of approximately 65-70 kDa, consistent with glycosylated form of CD86 (PMID: 8592066, 21849678).				
Notable Publications	Author	Pubmed ID Jour	nal	Application	
	Liping Xu	36179453 Tiss	ue Cell		
	Qiong Tang	36108436 Ecot	oxicol Environ Sa	f WB	
	Bo Zhang	32917854 Cell	Death Dis	WB	


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 13395-1-AP (CD86 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed raji cells using CD86 antibody (13395-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



1X10^6 Raji cells were stained with 0.2ug CD86 antibody (13395-1-AP, red) and control antibody (blue). Fixed with 90% MeOH blocked with 3% BSA (30 min). Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) with dilution 1:1000.