For Research Use Only SENP8 Polyclonal antibody

Catalog Number:13479-1-AP




Basic Information	Catalog Number: 13479-1-AP	GenBank Accession Number: BC031411	Purification Method: Antigen affinity purification	
	Size: 450 ug/ml Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG4287	GeneID (NCBI): 123228 UNIPROT ID: Q96LD8	Recommended Dilutions: IHC 1:250-1:1000 IF/ICC 1:200-1:800	
				Full Name: SUMO/sentrin specific peptidas family member 8
		Calculated MW: 212 aa, 24 kDa		
		Observed MW: 24-28 kDa		
		Applications	Tested Applications: IHC, IF/ICC, ELISA	Positive
Species Specificity: human, mouse, rat	colon ca			
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0			pancreas tissue, mouse kidney tissue IF/ICC : A431 cells,	
Background Information	SENP8, a human NEDD8-specific protease, also known as NEDP1. It is expected to be located in cytoplasm and nucleus, which is highly expressed in kidney and pancreas. The calculated molecular weight of SENP8 is 24 kDa. It is a protease that catalyzes two essential functions in the NEDD8 pathway: processing of full-length NEDD8 to its mature form and deconjugation of NEDD8 from targeted proteins such as cullins or p53 (PMID: 15242646). A single-residue difference in the C-terminus of NEDD8 and ubiquitin contributes significantly to the ability of NEDP1 to discriminate between them. In vivo analysis indicates that NEDP1 mutants perturb deNEDDylation of the tumour suppressor p53 (PMID: 12759363).			
Storage	Storage: Store at -20°C. Stable for one year Storage Buffer: PBS with 0.02% sodium azide and Aliquoting is unnecessary for -20°	50% glycerol pH 7.3.		


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 13479-1-AP (SENP8 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 13479-1-AP (SENP8 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human stomach cancer tissue slide using 13479-1-AP (SENP8 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded mouse pancreas tissue slide using 13479-1-AP (SENP8 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded rat kidney tissue slide using 13479-1-AP (SENP8 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded rat pancreas tissue slide using 13479-1-AP (SENP8 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded mouse kidney tissue slide using 13479-1-AP (SENP8 antibody) at dilution of 1:1 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse kidney tissue slide using 13479-1-AP (SENP8 antibody) at dilution of 1:1 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human liver tissue slide using 13479-1-AP (SENP8 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed A431 cells using SENP8 antibody (13479-1-AP) at dilution of 1:400 and Multi-rAb Coralite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002), CL594-phalloidin (red).