For Research Use Only

## P2RX4 Polyclonal antibody

Catalog Number:13534-1-AP

Featured Product

11 Publications





| Basic Information     | Catalog Number:<br>13534-1-AP  | GenBank Accession Number:<br>BC033826  | Purification Method:<br>Antigen affinity purification |  |
|-----------------------|--|--|---|--|
|                       | Size:  | GenelD (NCBI):   | Recommended Dilutions:                                |  |
|                       | 500 µg/ml<br>Source:   | 5025<br>UNIPROT ID:  | WB 1:1000-1:8000                                      |  |
|                       | Rabbit   | Q99571   |   |  |
|                       | Isotype:<br>IgG<br>Immunogen Catalog Number:<br>AG4445   | Full Name:<br>purinergic receptor P2X, ligand-gated<br>ion channel, 4  |   |  |
|                       |  | Calculated MW:<br>388 aa, 43 kDa   |   |  |
|                       |  | Observed MW:<br>45-60 kDa  |   |  |
| Applications          | Tested Applications:<br>WB, ELISA  | Positive Controls:<br>WB : HeLa cells,   |   |  |
|                       | Cited Applications:<br>IF, IHC, WB   |  |   |  |
|                       | Species Specificity:<br>human, mouse   |  |   |  |
|                       | Cited Species:<br>human, rat, mouse  |  |   |  |
| Background Informatio | extracellular nucleotides. These<br>that mediate calcium and potassi<br>receptors (GPCRs). P2RV4 (P2X pu<br>calculated molecular weight of P<br>reported, possibly due to post-tra | Nucleotides, the structural subunits of the nucleic acids, are also important extracellular signaling molecules. P2 receptors are a family of cell surface receptors that mediate a wide variety of physiologic effects in response to extracellular nucleotides. These receptors fall into two classes: P2X receptors, which are ligand-gated ion channels that mediate calcium and potassium fluxes in response to ATP, and P2Y receptors, which are G protein-coupled receptors (GPCRs). P2RX4 (P2X purinoceptor 4) is a receptor for ATP acting as a ligand gated ion channel. The calculated molecular weight of P2RX4 is 43 kDa, larger apparent molecular weights of 50-80 kDa have been reported, possibly due to post-translational glycosylation (PMID: 24040145, 15262999; 29382907; 12566439). The antibody can detect the band around 45 kDa. |   |  |
| Notable Publications  | Author   | Pubmed ID Journal  | Application   |  |
|                       | Yexian Yuan  | 35507647 Sci Adv   | WB  |  |
|                       | Si-Ting Wu   | 35352488 CNS Neurosci The  | r IF  |  |
|                       | Zhuo Lin   | 33828988 Front Oncol   | IHC   |  |
| Storage               | Storage:<br>Store at -20°C. Stable for one yea<br>Storage Buffer:<br>PBS with 0.02% sodium azide and   |  |   |  |

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data





WB result of P2RX4 antibody (13534-1-AP; 1:4000; room temperature for 1.5 hours) with sh-control and sh-P2RX4 transfected HeLa cells. WB result of P2RX4 antibody (13534-1-AP; 1:4000; room temperature for 1.5 hours) with wild-type and P2RX4 knockout HeLa cells.