

For Research Use Only

PGK2 Polyclonal antibody

Catalog Number: 13686-1-AP

Featured Product

5 Publications



Basic Information

Catalog Number:

13686-1-AP

Size:

400 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG4553

GenBank Accession Number:

BC038843

GeneID (NCBI):

5232

UNIPROT ID:

P07205

Full Name:

phosphoglycerate kinase 2

Calculated MW:

417 aa, 45 kDa

Observed MW:

45 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:1000-1:4000

IHC 1:100-1:400

IF-P 1:50-1:500

Applications

Tested Applications:

WB, IHC, IF-P, ELISA

Cited Applications:

WB

Species Specificity:

human, mouse

Cited Species:

human, mouse, sheep

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : mouse testis tissue, HeLa cells, HepG2 cells, human heart tissue, human testis tissue, mouse brain tissue

IHC : human testis tissue,

IF-P: mouse testis tissue,

Background Information

PGK2, phosphoglycerate kinase 2, which belongs to the phosphoglycerate kinase superfamily. It catalyzes the first ATP-generating step in the central metabolic pathway of glycolysis, converting 1,3-bisphosphoglycerate and ADP to 3-phosphoglycerate and ATP.

Notable Publications

Author	Pubmed ID	Journal	Application
Guangjian Jiang	27832638	Cell Physiol Biochem	WB
Xiaolei Yao	28460680	Theriogenology	WB
Huilin Li	36517592	Nature	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




mouse testis tissue were subjected to SDS PAGE followed by western blot with 13686-1-AP (PGK2 antibody) at dilution of 1:1200 incubated at room temperature for 1.5 hours.




Immunofluorescent analysis of (4% PFA) fixed mouse testis tissue using PGK2 antibody (13686-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed mouse testis tissue using PGK2 antibody (13686-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunohistochemical analysis of paraffin-embedded human testis tissue slide using 13686-1-AP (PGK2 Antibody) at dilution of 1:200 (under 10x lens).



Immunohistochemical analysis of paraffin-embedded human testis tissue slide using 13686-1-AP (PGK2 Antibody) at dilution of 1:200 (under 40x lens).