

For Research Use Only

FGF12 Polyclonal antibody

Catalog Number: 13784-1-AP

Featured Product

4 Publications



Basic Information

Catalog Number:

13784-1-AP

Size:

260 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG4797

GenBank Accession Number:

BC022524

GeneID (NCBI):

2257

UNIPROT ID:

P61328

Full Name:

fibroblast growth factor 12

Calculated MW:

181 aa, 20 kDa

Observed MW:

20-22 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:2000-1:12000

IP 0.5-4.0 µg for 1.0-3.0 mg of total protein lysate

IHC 1:20-1:200

Applications

Tested Applications:

IHC, IP, WB, ELISA

Cited Applications:

WB, IHC, IF

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : mouse brain tissue, rat brain tissue

IP : mouse brain tissue,

IHC : mouse brain tissue,

Background Information

FGF12 is a member of the FGF family. FGF family members play important roles in embryogenesis, angiogenesis, and wound repair. FGF12 lacks the N-terminal signal sequence present in most of the FGF family members, but it contains clusters of basic residues that have been demonstrated to act as a nuclear localization signal. When transfected into mammalian cells, this protein accumulated in the nucleus, but was not secreted. FGF12 plays an intracellular role in the inhibition of radiation-induced apoptosis. FGF12 is expressed abundantly in developing and adult nervous systems; therefore, FGF12 was thought to be related to nervous system development and function.

Notable Publications

| Author | Pubmed ID | Journal | Application |
|----------------|-----------|-----------|-------------|
| Jana Velišková | 33982289 | Epilepsia | WB |
| Yukiko Hanada | 30065296 | Sci Rep | WB,IF |
| Ya Zhang | 39091933 | Heliyon | IHC |

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 13784-1-AP (FGF12 antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.



IP result of anti-FGF12 (IP:13784-1-AP, 4ug; Detection:13784-1-AP 1:6000) with mouse brain tissue lysate 1280 ug.



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 13784-1-AP (FGF12 Antibody) at dilution of 1:50 (under 10x lens).



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 13784-1-AP (FGF12 Antibody) at dilution of 1:50 (under 40x lens).