For Research Use Only

SEC63 Polyclonal antibody

Catalog Number: 13978-1-AP

Featured Product

5 Publications



Basic Information

Catalog Number: GenBank Accession Number: 13978-1-AP BC047221

Size: GeneID (NCBI): 11231

Source: UNIPROT ID: Rabbit Q9UGP8

Isotype: Full Name:

gG SEC63 homolog (S. cerevisiae)

Immunogen Catalog Number:Calculated MW:AG502188 kDa

Observed MW: 83-90 kDa Antigen affinity purification Recommended Dilutions: WB 1:1000-1:4000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:50-1:500 IF-P 1:50-1:500 IF/ICC 1:50-1:500

Purification Method:

Applications

Tested Applications:

WB, IHC, IF/ICC, IF-P, IP, ELISA

Cited Applications:

WB, IF

Species Specificity: human, mouse, rat Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: A431 cells, HEK-293 cells, PC-3 cells

IP: A431 cells,

IHC: human liver tissue, mouse pancreas tissue

IF-P: mouse kidney tissue, IF/ICC: HEK-293 cells,

Background Information

SEC63 is an integral ER membrane protein. SEC63 and SEC62 form a dimeric complex and play a central role in the translocation of nascent and newly synthesized precursor polypeptides into the ER. Mutations in the gene encoding SEC63 cause polycystic liver disease in humans. The intact SEC63 protein is usually detected around 83-90 kDa, while smaller cleavage products can be observed by autophagy proteolysis (PMID: 25052096)

Notable Publications

Author	Pubmed ID	Journal	Application
Tetsuya Hirata	36055406	J Biol Chem	WB
Hossam Hassan	26303067	J Am Soc Nephrol	WB, IF
Brian J Conti	25801167	Mol Cell	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




A431 cells were subjected to SDS PAGE followed by western blot with 13978-1-AP (SEC 63 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours




Immunofluorescent analysis of (4% PFA) fixed mouse kidney tissue using 13978-1-AP (SEC63 antibody) at dilution of 1:50 and Alexa Fluor 488-Conjugated Goat Anti-Rabbit IgG(H+L).




Immunofluorescent analysis of (4% PFA) fixed mouse kidney tissue using 13978-1-AP (SEC63 antibody) at dilution of 1:50 and Alexa Fluor 488-Conjugated Goat Anti-Rabbit IgG(H+L).




Immunohistochemical analysis of paraffinembedded human liver tissue slide using 13978-1-AP (SEC63 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human liver tissue slide using 13978-1-AP (SEC63 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-SEC 63 (IP:13978-1-AP, 4ug; Detection:13978-1-AP 1:300) with A431 cells lysate 2400ug.



Immunofluorescent analysis of (-20°C Ethanol) fixed HEK-293 cells using SEC63 antibody (13978-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L).