

For Research Use Only

OLIG2 Polyclonal antibody

Catalog Number: 13999-1-AP **60 Publications**



Basic Information

Catalog Number:

13999-1-AP

Size:

500 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG5089

GenBank Accession Number:

BC047511

GeneID (NCBI):

10215

UNIPROT ID:

Q13516

Full Name:

oligodendrocyte lineage transcription factor 2

Calculated MW:

32 kDa

Observed MW:

32-36 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:1000-1:8000

IP 0.5-4.0 µg for 1.0-3.0 mg of total protein lysate

IHC 1:50-1:500

IF 1:250-1:1000

Applications

Tested Applications:

IF-P, IHC, IP, WB, ELISA

Cited Applications:

IF, IHC, WB

Species Specificity:

human, mouse, rat

Cited Species:

human, rat, mouse, bovine

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : mouse brain tissue, rat brain

IP : mouse brain tissue,

IHC : mouse brain tissue, human gliomas tissue

IF : rat brain tissue,

Background Information

OLIG2, also named as BHLHB1, BHLHE19, PRKCBP2 and RACK17, is required for oligodendrocyte and motor neuron specification in the spinal cord, as well as for the development of somatic motor neurons in the hindbrain. Cooperates with OLIG1, OLIG2 establish the pMN domain of the embryonic neural tube. Antagonist of V2 interneuron and of NIKX2-2-induced V3 interneuron development. OLIG2 is widely expressed in subsets of glia cells and progenitors, and it is strongly induced at different sites by both acute and chronic injury, albeit with different mechanisms. OLIG2 acts as a repressor of neurogenesis in cells reacting to brain injury. It may represent an effective approach towards evoking neuronal repair from parenchymal precursors.(PMID:19390819)

Notable Publications

Author	Pubmed ID	Journal	Application
Lirong Liang	34585785	J Pineal Res	WB
Yizi Zhu	36142668	Int J Mol Sci	IF
Angela M Lager	30213958	Nat Commun	IF

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 13999-1-AP (OLIG2 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 13999-1-AP (OLIG2 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 13999-1-AP (OLIG2 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed rat brain tissue using 13999-1-AP (OLIG2 antibody), at dilution of 1:500 and CoralLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



IP result of anti-OLIG2 (IP:13999-1-AP, 4ug; Detection:13999-1-AP 1:1000) with mouse brain tissue lysate 1600 ug.