### For Research Use Only

# USP16 Polyclonal antibody

Catalog Number: 14055-1-AP

Featured Product

5 Publications



**Basic Information** 

Catalog Number:

GenBank Accession Number:

14055-1-AP

BC030777

Size:

GeneID (NCBI):

500 ug/ml

10600

Source:

UNIPROT ID:

Rabbit

Q9Y5T5

Isotype:

Full Name:

IgG ubiquitin specific peptidase 16

Immunogen Catalog Number: Calculated MW: AG4981 94 kDa

Observed MW: 100-120 kDa Purification Method: Antigen affinity purification Recommended Dilutions:

WB 1:500-1:2000

IHC 1:50-1:500

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

**Applications** 

Tested Applications: WB, IHC, IP, ELISA Cited Applications: WB, IHC, IF, IP, CoIP Species Specificity: human

**Cited Species:** 

human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HeLa cells, K-562 cells

IP: HeLa cells.

IHC: human liver tissue, human brain tissue, human spleen tissue, human ovary tissue, human kidney tissue, human heart tissue, human testis tissue, human skin tissue

# **Background Information**

USP16 (Ubiquitin specific peptidase 16) was originally identified as a DUB of histone H2A which is implicated in cell cycle progression and HOXD10 gene expression (PMID:17914355). The knockout study has reported that depletion of USP16 results in early embryonic lethality in mice (PMID:24784029). Moreover, USP16 mediated the deubiquitination and stabilization of Drp1 through its direct interaction with Drp1 (PMID:37488647).

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Christine Galant	30608617	Rheumatology (Oxford)	IHC
Nadine Sen Nkwe	34005181	J Cell Sci	WB,IF
Nadine Sen Nkwe	32005696	J Cell Sci	WB,IF

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**



Various lysates were subjected to SDS PAGE followed by western blot with 14055-1-AP (USP16 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 14055-1-AP (USP16 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 14055-1-AP (USP16 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human liver tissue slide using 14055-1-AP (USP16 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human liver tissue slide using 14055-1-AP (USP16 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-USP16 (IP:14055-1-AP, 5ug; Detection:14055-1-AP 1:1000) with HeLa cells lysate 880ug.