## For Research Use Only

# **HVCN1** Polyclonal antibody

Catalog Number:14162-1-AP 1 Publications



**Basic Information** 

Catalog Number: 14162-1-AP

GenBank Accession Number: BC032672

**Purification Method:** Antigen affinity purification

Size:

GeneID (NCBI): 84329

Recommended Dilutions:

350 ug/ml Source:

**UNIPROT ID:** 

WB 1:500-1:1000 IP 0.5-4.0 ug for 1.0-3.0 mg of total

Rabbit

Q96D96

Isotype:

Full Name:

protein lysate

hydrogen voltage-gated channel 1 Calculated MW:

IHC 1:50-1:500 IF/ICC 1:20-1:200

Immunogen Catalog Number:

AG5350 273 aa, 32 kDa

Observed MW:

28-32 kDa, ~60 kDa

**Applications** 

**Tested Applications:** 

WB, IHC, IF/ICC, IP, ELISA

WB: Raji cells, PC-3 cells

**Cited Applications:** 

IP: PC-3 cells,

IHC: human tonsillitis tissue,

Species Specificity: human

IF/ICC: PC-3 cells,

**Positive Controls:** 

Cited Species:

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval antigen retrieval with a trace buffer and 16.0 with citrate buffer pH 6.0

# **Background Information**

HVCN1, also named as VSOP and HV1, Belongs to the hydrogen channel family. HVCN1 mediates the voltagedependent proton permeability of excitable membranes. It forms a proton-selective channel through which protons may pass in accordance with their electrochemical gradient. Proton efflux, HVCN1 is accompanied by membrane depolarization, facilitates acute production of reactive oxygen species in phagocytosis. HVCN1, the voltage $sensitive\ proton\ channel, is\ present\ in\ human\ sperm\ and\ is\ an\ important\ regulator\ of\ the\ functional\ maturation\ of\ present\ in\ pr$ sperm. HVCN1 has four isoforms with MW 28-32 kDa or 40 kDa (modification). It has a dimer form with MW ~60 kDa.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Bing Yu	33936061	Front Immunol	IHC

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

## **Selected Validation Data**




IP result of anti-HVCN1 (IP:14162-1-AP, 4ug; Detection:14162-1-AP 1:300) with PC-3 cells lysate




Raji cells were subjected to SDS PAGE followed by western blot with 14162-1-AP (HVCN1 Antibody) at dilution of 1:600 incubated at room temperature for 15 hours.



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 14162-1-AP (HVCN1 antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed PC-3 cells using 14162-1-AP (HVCN1 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 14162-1-AP (HVCN1 Antibody) at dilution of 1:200 (under 40x lens).