## For Research Use Only

## HLX Polyclonal antibody

Catalog Number: 14336-1-AP

**Featured Product** 

1 Publications

BC033808

GeneID (NCBI):

**UNIPROT ID:** 

Q14774 Full Name:

51 kDa

GenBank Accession Number:



**Basic Information** 

Catalog Number: 14336-1-AP

Size: 350 μg/ml Source: Rabbit Isotype:

Immunogen Catalog Number:

AG5627

H2.0-like homeobox Calculated MW: 51 kDa Observed MW:

**Purification Method:** 

Antigen affinity purification Recommended Dilutions:

WB 1:500-1:1000 IHC 1:20-1:200 IF/ICC 1:50-1:500

**Applications** 

**Tested Applications:** WB, IHC, IF/ICC, ELISA Cited Applications:

WB, IHC

Species Specificity:

human **Cited Species:** human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

**Positive Controls:** 

WB: human placenta tissue,

IHC: human endometrial cancer tissue,

IF/ICC: A549 cells,

**Background Information** 

In immune system, HIx is a Th1-special transcription factor that interacts specifically with T-bet, and synergistically they promote IFN- γ expression when co-expressed. It can down-regulate the IL-4 receptor expression in naive CD4+ T cells [PMID:14688316]. Over expression of HIx contributes ot the aberrant expression of IFN- $\gamma$  in normal CD4+T cells during the differentiation under Th2-polarizing conditions. In addition, Hlx is a negative regulator of IFN-  $\gamma$ production in NK cells and its inhibitory function is achieved at least in part through the proteasomal degradation of STAT4 [PMID:16424184].

**Notable Publications** 

Author **Pubmed ID** Journal Application Ting Liu 26631039 **Tumour Biol** WB,IHC


Storage

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data




Immunofluorescent analysis of (4% PFA) fixed A549 cells using 14336-1-AP (HLX antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



human placenta tissue were subjected to SDS PAGE followed by western blot with 14336-1-AP (HLX antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human endometrial cancer using 14336-1-AP (HLX antibody) at dilution of 1:100 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human endometrial cancer using 14336-1-AP (HLX antibody) at dilution of 1:100 (under 40x lens).