## For Research Use Only

# NDUFB8 Polyclonal antibody

Catalog Number: 14794-1-AP

Featured Product

68 Publications



**Basic Information** 

Catalog Number: 14794-1-AP Source: Rabbit

Isotype: IgG

Immunogen Catalog Number:

AG6495

GenBank Accession Number:

BC000466
GeneID (NCBI):
4714
UNIPROT ID:

095169

Full Name: NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 8, 19kDa

Calculated MW: 22 kDa

Observed MW: 19-22 kDa

Applications

Tested Applications: WB, IHC, IF/ICC, IP, ELISA Cited Applications: WB, IHC, IF

Species Specificity: human, mouse, rat Cited Species:

human, mouse, rat, pig, canine, zebrafish

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 **Positive Controls:** 

WB: HepG2 cells, A549 cells, various samples, 3T3-L1 cells, mouse brain tissue, mouse liver tissue, rat brain

**Purification Method:** 

WB: 1:5000-1:50000

IF/ICC: 1:200-1:800

protein lysate

IHC: 1:50-1:500

Antigen affinity purification

IP: 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

IP: HepG2 cells,

IHC: human liver tissue, human heart tissue, mouse

brain tissue

IF/ICC: HepG2 cells,

# **Background Information**

## Notable Publications

Author	Pubmed ID	Journal	Application
Marine C N M Blackman	36139533	Cancers (Basel)	WB
Delong Yin	34497154	Aging (Albany NY)	WB
Kazuhide Takahashi	28890359	Exp Gerontol	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol, pH7.3  $\,$ 

Aliquoting is unnecessary for -20°C storage


#### Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 14.794-1-AP (NDUFB8 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human liver tissue slide using 14794-1-AP (NDUFB8 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with PK30006 (OXPHOS Cocktail (Human recommended)) at dilution of 1:4000 incubated at room temperature for 1 hours.




IP result of anti-NDUFB8 (IP:14794-1-AP, 4ug; Detection:14794-1-AP 1:2000) with HepG2 cells lysate 880 ug.



WB result of NDUFB8 antibody (14794-1-AP; 1:40000; incubated at room temperature for 1.5 hours) with sh-Control and sh-NDUFB8 transfected A549 cells.



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 14794-1-AP (NDUFB8 antibody) at dilution of 1:800 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using NDUFB8 antibody (14794-1-AP) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).