For Research Use Only

Hikeshi Polyclonal antibody

Catalog Number:14808-1-AP

Featured Product

5 Publications

22 kDa



Basic Information

14808-1-AP Size: 187 µg/ml Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG6530

Catalog Number:

GenBank Accession Number: BC001677 GeneID (NCBI): 51501 UNIPROT ID: Q53FT3 Full Name: chromosome 11 open reading frame 73 Calculated MW: 22 kDa Observed MW:

Purification Method: Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:1000 IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate IHC 1:20-1:200

Applications

Tested Applications: IHC, IP, WB,ELISA Cited Applications: WB, IHC Species Specificity: human, mouse, rat Cited Species:

human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : mouse kidney tissue, A375 cells, mouse brain tissue, PC-3 cells IP : mouse brain tissue, IHC : human kidney tissue,

Background Information

Notable Publications	Author	Pubmed ID	Journal	Application
	K Berthenet	27819670	Oncogene	
	Simon Edvardson	26545878	J Med Genet	WB
	Antonios Apostolopoulos	30619453	Front Genet	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment. Storage Buffer: PBS with 0.02% sodium azide and 50% glycerol pH 7.3. Aliquoting is unnecessary for -20°C storage


 For technical support and original validation data for this product please contact:


 T: 4006900926
 E: Proteintech-CN@ptglab.com
 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data








mouse kidney tissue were subjected to SDS PAGE followed by western blot with 14808-1-AP (Hikeshi antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.

Immunohistochemical analysis of paraffinembedded human kidney using 14808-1-AP (Hikeshi antibody) at dilution of 1:50 (under 10x lens). Immunohistochemical analysis of paraffinembedded human kidney using 14808-1-AP (Hikeshi antibody) at dilution of 1:50 (under 40x lens).



IP result of anti-Hikeshi (IP:14808-1-AP, 4ug; Detection:14808-1-AP 1:500) with mouse brain tissue lysate 2640ug.