For Research Use Only

DAAM1 Polyclonal antibody

Catalog Number: 14876-1-AP

Featured Product

22 Publications



Basic Information

Catalog Number:

GenBank Accession Number:

14876-1-AP

BC064999

Size:

GeneID (NCBI):

23002

Source:

UNIPROT ID:

Rabbit

Q9Y4D1

Isotype:

Full Name:

IgG dishevelled associated activator of morphogenesis 1

AG6648 Calculated MW: 123 kDa
Observed MW:

123 kDa

Purification Method:

Antigen affinity purification Recommended Dilutions:

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

protein lysate IHC 1:20-1:200 IF 1:20-1:200

WB 1:1000-1:4000

Applications

Tested Applications: IF/ICC, IHC, IP, WB,ELISA Cited Applications: WB, IF, IHC, CoIP Species Specificity: human, mouse, rat

Cited Species:

human, goat, mouse, Xenopus, Canine

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: mouse brain tissue, HepG2 cells, human heart tissue, mouse liver tissue, mouse heart tissue

IP: mouse liver tissue,
IHC: human brain tissue,
IF: HepG2 cells,

Background Information

Dishevelled-associated activator of morphogenesis 1 (Daam1) is a member of the formin protein family that is implicated in actin assembly. Daam1 also plays a critical role in mediating Wnt/Fz signaling pathway and regulates cell polarity during development and tissue homoeostasis. This antibody well recognizes the endogenous DAAM1 protein around 120 kDa.

Notable Publications

Author	Pubmed ID	Journal	Application
Qin Ju He	31618077	Am J Physiol Cell Physiol	WB
Tamako Nishimura	27807130	J Cell Biol	WB,IF
Jie Mei	31757662	Pathol Res Pract	WB,IHC

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



mouse brain tissue were subjected to SDS PAGE followed by western blot with 14876-1-AP (DAAM1 Antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.




Immunofluorescent analysis of HepG2 cells, using DAAM1 antibody 14876-1-AP at 1:50 dilution and Rhodamine-labeled goat anti-rabbit IgG (red). Blue pseudocolor = DAPI (fluorescent DNA dye).



Immunohistochemical analysis of paraffinembedded human brain using 14876-1-AP (DAAM1 antibody) at dilution of 1:100 (under 40x lens).



Immunohistochemical analysis of paraffinembedded human brain using 14876-1-AP (DAAM1 antibody) at dilution of 1:100 (under 10x lens).



IP result of anti-DAAM1 (IP:14876-1-AP, 3ug; Detection:14876-1-AP 1:500) with mouse liver tissue lysate 8000ug.