For Research Use Only

NGRN Polyclonal antibody

Catalog Number: 14885-1-AP

Featured Product

4 Publications



Basic Information

 Catalog Number:
 GenBank Accession Number:

 14885-1-AP
 BC001682

 Size:
 GeneID (NCBI):

 1000 μ g/ml
 51335

 Source:
 UNIPROT ID:

Rabbit Q9NPE2 Isotype: Full Name:

IgG neugrin, neurite outgrowth associated

Immunogen Catalog Number: Calculated MW:

AG6661 32 kDa

Observed MW: 30-32 kDa

Applications

Tested Applications: IHC, WB, ELISA Cited Applications:

M/D IE

WB, IF

Species Specificity: human, mouse, rat Cited Species: human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: mouse liver tissue,

 $IHC: human\, brain\, tissue,\, human\, heart\, tissue,\, human\,$

Purification Method:

WB 1:200-1:1000 IHC 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

testis tissue

Background Information

NGRN, also named as Neugrin or F158G, is a 291 amino acid protein, which belongs to the neugrin family. NGRN localizes in the nucleus and is expressed at high levels in heart, brain and skeletal muscle. In brain, it is mainly expressed in neurons rather than glial cells. NGRN may be involved in neuronal differentiation.

Notable Publications

Author	Pubmed ID	Journal	Application
Jason D Arroyo	27667664	Cell Metab	WB,IF
Víctor Llombart	27888142	J Proteomics	
Hana Antonicka	27974379	EMBO Rep	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

Selected Validation Data

Immunohistochemical analysis of paraffinembedded human brain tissue slide using 14885-1-AP (NGRN Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human brain tissue slide using 14885-1-AP (NGRN Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



mouse liver tissue were subjected to SDS PAGE followed by western blot with 14885-1-AP (NGRN Antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.