For Research Use Only

IL-10RB Polyclonal antibody

Catalog Number:15102-1-AP

3 Publications



Basic Information

Applications

Catalog Number: 15102-1-AP

Size: 450 µg/ml Source: Rabbit Isotype:

Immunogen Catalog Number:

AG7445

Observed MW: 42-45 kDa

GenBank Accession Number:

interleukin 10 receptor, beta

BC001903

3588

GeneID (NCBI):

UNIPROT ID:

Calculated MW:

Q08334 Full Name:

37 kDa

Antigen affinity purification Recommended Dilutions: WB 1:300-1:1000

Purification Method:

Tested Applications:

WB, ELISA

Cited Applications:

WB

Species Specificity: human, mouse, rat Cited Species: human, mouse

WB: rat skeletal muscle tissue, Jurkat cells, mouse skeletal muscle tissue

Background Information

Interleukin-10 (IL-10) is an anti-inflammatory cytokine that plays a critical role in maintaining immune system balance. IL-10 signals through a receptor complex consisting of two molecules of the IL-10R alpha-chain (IL-10RA) and two molecules of the accessory IL-10R beta-chain (IL-10RB) (PMID: 22236434). IL-10RA and IL-10RB are members of the type II cytokine receptor family. IL-10RB, also known as IL-10R2 or CDw210b, is the signaling subunit of the IL-10R complex and is constitutively expressed in a variety of cell types (PMID: 24507158). Binding of IL-10 to the IL-10R complex leads to JAK1- and TYK2-mediated phosphorylation of STAT3 (PMID: 11244051; 10433356). IL-10RB is also shared by several other cytokines, including IL22, IL-26, IFN- γ, IL-28A/B and IL29 (PMID: 22236434).

Notable Publications

Author	Pubmed ID	Journal	Application
Yang Che	34015450	Biochim Biophys Acta Mol Basis Dis	WB
Shuyue Luo	35811529	J Alzheimers Dis	WB
Mai Yamamoto	37319277	PLoS One	WB


Storage

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



rat skeletal muscle tissue were subjected to SDS PAGE followed by western blot with 15102-1-AP (IL10RB antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.