

For Research Use Only

MVD Polyclonal antibody

Catalog Number: 15331-1-AP **4 Publications**



Basic Information

Catalog Number: 15331-1-AP	GenBank Accession Number: BC000011	Purification Method: Antigen affinity purification
Size: 240 µg/ml	GeneID (NCBI): 4597	Recommended Dilutions: WB 1:500-1:2000 IHC 1:50-1:500 IF/ICC 1:50-1:500
Source: Rabbit	UNIPROT ID: P53602	
Isotype: IgG	Full Name: mevalonate (diphospho) decarboxylase	
Immunogen Catalog Number: AG7255	Calculated MW: 43 kDa	
	Observed MW: 66-74 kDa, 45 kDa, 37 kDa	

Applications

Tested Applications: WB, IHC, IF/ICC, ELISA	Positive Controls: WB : HCT 116 cells, rat liver tissue, HepG2 cells, K-562 cells IHC : human colon tissue, human lung cancer tissue, human heart tissue IF/ICC : A431 cells,
Cited Applications: WB, IHC	
Species Specificity: human, mouse, rat	
Cited Species: human, mouse	

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

The enzyme mevalonate pyrophosphate decarboxylase(MVD) catalyzes the conversion of mevalonate pyrophosphate into isopentenyl pyrophosphate. It is also named as MPD and as a unique enzyme in one of the early steps in cholesterol biosynthesis, MVD may be a useful target for drugs aimed at lowering serum cholesterol levels(PMID:8626466). The intracellular glycosylation does not contribute to the difference between the 45 and 37 kDa species of MVD. The native MVD has a molecular weight of 90 kDa that it consists of two identical subunits of 45 kDa and a 37 kDa protein is also found as a subunit of MVD and this type of MVD may be a 74 kDa. But the 37 kDa enzyme appeared only when the rats are fed the CP diet.(PMID:9348097).

Notable Publications

Author	Pubmed ID	Journal	Application
Zhenhua Zhang	34562605	Cell Signal	IHC
Audrey Basque	35723385	Curr Issues Mol Biol	WB
Kailin Xing	38084209	J Hepatocell Carcinoma	WB

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




HepG2 cells were subjected to SDS PAGE followed by western blot with 15331-1-AP (MVD antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.




Various lysates were subjected to SDS PAGE followed by western blot with 15331-1-AP (MVD antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human lung cancer tissue slide using 15331-1-AP (MVD Antibody) at dilution of 1:50 (under 10x lens).



Immunohistochemical analysis of paraffin-embedded human lung cancer tissue slide using 15331-1-AP (MVD Antibody) at dilution of 1:50 (under 40x lens).



Immunohistochemical analysis of paraffin-embedded human colon tissue slide using 15331-1-AP (MVD antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (10% Formaldehyde) fixed A431 cells using 15331-1-AP (MVD antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).