For Research Use Only

Beta Galactosidase Polyclonal antibody, PBS Only

Catalog Number: 15518-1-PBS



Purification Method:

Antigen affinity purification

Basic Information

Catalog Number: 15518-1-PBS

BC007493 GeneID (NCBI):

GenBank Accession Number:

 Concentration:
 GeneID (NCE)

 1 mg/ml
 2720

 Source:
 UNIPROT ID:

 Rabbit
 P16278

Isotype: Full Name:
IgG galactosidase, beta 1

7792 76 kDa

Observed MW: 67 kDa

Calculated MW:

Applications

Tested Applications: WB, IHC, IF/ICC, IF-P, ELISA Species Specificity:

Immunogen Catalog Number:

human, mouse

Background Information

GLB1(Beta-galactosidase) is also named as ELNR1 or Lactase. It cleaves beta-linked terminal galactosyl residues from gangliosides, glycoproteins, and glycosaminoglycans. This protein is identical to the elastin-binding protein (EBP), a major component of the nonintegrin cell surface receptor complex expressed in fibroblasts, smooth muscle cells, chondroblasts, leukocytes, and certain cancer cell types. Defects in GLB1 are the cause of GM1-gangliosidosis type 1 (GM1G1), GM1-gangliosidosis type 2 (GM1G2), GM1-gangliosidosis type 3 (GM1G3) and mucopolysaccharidosis type 4B (MPS4B). GBL1 is synthesized as an 85-kDa precursor that is C-terminally processed into a 64-66 kDa mature form and the released ~20-kDa proteolytic fragment was thought to be degraded (PMID: 10744681). GLB1 has 3 isoforms with MW of 76 kDa, 73 kda and 61 kDa.

Storage

Storage:

Store at -80°C.


The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer: PBS only, pH7.3


Selected Validation Data




SH-SY5Y cells were subjected to SDS PAGE followed by western blot with 15518-1-AP (GLB1 antibody) at dilution of 1:400 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 15518-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human prostate cancer tissue slide using 15518-1-AP (Beta Galactosidase antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 15518-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human prostate cancer tissue slide using 15518-1-AP (Beta Galactosidase antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 15518-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using Beta Galactosidase antibody (15518-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red). This data was developed using the same antibody clone with 15518-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded human prostate cancer tissue using Beta Galactosidase antibody (15518-1-AP) at dilution of 1:200 and Multi-Ab Coralite @ Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 15518-1-PBS in a different storage buffer



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded human liver cancer tissue using Beta Galactosidase antibody (15518-1-AP) at dilution of 1:200 and Multi-rAb Coralite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 15518-1-PBS in a different storage buffer formulation.