For Research Use Only

ARHGDIB Polyclonal antibody

Catalog Number: 16122-1-AP

Featured Product

6 Publications

BC009200

GenBank Accession Number:



Basic Information

Catalog Number: 16122-1-AP

Size: GeneID (NCBI): 1000 μ g/ml 397

Source: UNIPROT ID: Rabbit P52566
Isotype: Full Name:

gG Rho GDP dissociation inhibitor (GDI)

Immunogen Catalog Number: b

AG9078 Calculated MW: 201 aa, 23 kDa

Observed MW: 27 kDa

Applications

Tested Applications: IHC, IP, WB, ELISA Cited Applications:

WB, IHC

Species Specificity: human, mouse, rat Cited Species:

human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 **Positive Controls:**

WB: HL-60 cells, K-562 cells, rat spleen tissue, Jurkat

Purification Method:

WB 1:5000-1:50000

protein lysate

IHC 1:20-1:200

Antigen affinity purification

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

cells, Raji cells IP: Raji cells,

IHC: human stomach cancer tissue,

Background Information

ARHGDIB, also known as RhoGDI2, is a conserved member of the RhoGDI family and plays an important role in cell migrations. It regulates the GDP/GTP exchange reaction of the Rho proteins by inhibiting the dissociation of GDP from them, and the subsequent binding of GTP to them. It is mainly in hematopoietic, endothelial, and epithelial cells. It has been linked to tumorigenesis and metastasis. RhoGDI2 expression is downregulated in several cancer types, such as bladder, lung and lymphoma, but is upregulated in prostate and gastric cancer.

Notable Publications

Author	Pubmed ID	Journal	Application
Yang Fang	25180836	DNA Cell Biol	WB
Fan Zhang	25411139	Proteomics	WB
Lei Liu	34006303	Mol Cancer	WB

Storage

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 16122-1-AP (ARHGDIB antibody) at dilution of 1:40000 incubated at room temperature for 1.5 hours.



IP result of anti-ARHGDIB (IP:16122-1-AP, 4ug; Detection:16122-1-AP 1:1000) with Raji cells lysate 2000ug.



Immunohistochemical analysis of paraffinembedded human stomach cancer using 16122-1-AP (ARHGDIB antibody) at dilution of 1:100 (under 10x Jens)



Immunohistochemical analysis of paraffinembedded human stomach cancer using 16122-1-AP (ARHGDIB antibody) at dilution of 1:100 (under 40x lens).