For Research Use Only

NXN Polyclonal antibody

Catalog Number:16128-1-AP

Featured Product

7 Publications




Basic Information	Catalog Number: 16128-1-AP	GenBank Accession Number: BC009327		cation Method: en affinity purification	
	Size:	GenelD (NCBI):	-	mmended Dilutions:	
	350 µg/ml	64359		:1000-1:6000	
	Source: UNIPROT ID: Rabbit Q6DKJ4			IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate	
	lsotype: lgG	Full Name: nucleoredoxin		IHC 1:50-1:500 IF 1:50-1:500	
	Immunogen Catalog Number: AG9113	Calculated MW: 435 aa, 48 kDa			
		Observed MW: 48 kDa			
Applications	Tested Applications:		Positive Controls:		
	IF/ICC, IHC, IP, WB, ELISA			eLa cells, human heart tissue, H	
	Cited Applications: WB, IF, IHC	WB IE HC		ells, HepG2 cells, SH-SY5Y cells	
	Species Specificity:		IP : HeLa cells, IHC : human colon c	ancertissue	
	human, mouse, rat		IF : HeLa cells,		
	Cited Species: In The Sector, human, mouse, rat				
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0				
Background Information	Nucleoredoxin (NRX/NXN), a relat reported to regulate the Wnt/β-c dependent manner(PMID:175672 mouse tissues, with highest expr isoforms produced by alternative	catenin pathway, which itse (40). Northern blot analysis ession in testis and skin an	elf regulates cell fate a detected several nucl d lowest expression ir	and early development, in a red eoredoxin splice variants in adu	
Notable Publications	Author	Pubmed ID Journ	al	Application	
	Claudia Urbainsky	30584462 Oxid	Med Cell Longev	WB	
	Kendra M Prutton	35537597 Free	Radic Biol Med	WB	
	Vedangi Hambardikar	35453370 Antic	xidants (Basel)	WB	
	Vedangi Hambardikar	35453370 Antic	xidants (Basel)	WB	


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 16128-1-AP (NXN antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using NXN antibody (16128-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594phalloidin (red).



IP result of anti-NXN (IP:16128-1-AP, 4ug; Detection:16128-1-AP 1:500) with HeLa cells lysate 3200ug.



Various lysates were subjected to SDS PAGE followed by western blot with 16128-1-AP (NXN antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.