For Research Use Only

PSEN2-Specific Polyclonal antibody

Catalog Number: 16168-1-AP

Featured Product

2 Publications



Basic Information

Catalog Number: 16168-1-AP

Size: 650 µg/ml Source: Rabbit

Isotype: IgG GenBank Accession Number:

NM_000447 GeneID (NCBI): 5664 UNIPROT ID: P49810

presenilin 2 (Alzheimer disease 4)

Calculated MW: 50 kDa Observed MW: 55-60 kDa

Full Name:

Purification Method:

Antigen affinity purification Recommended Dilutions:

WB 1:300-1:800 IHC 1:20-1:200

Applications

Tested Applications: IHC, WB, ELISA Cited Applications:

WB

Species Specificity: human, mouse, rat Cited Species: mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: C6 cells, A549 cells, NIH/3T3 cells

IHC: human kidney tissue, human placenta tissue, human colon tissue, mouse brain tissue, human breast cancer tissue

Background Information

Presenilins can regulate APP processing through their effects on gamma-secretase, an enzyme that cleaves APP. These transmembrane proteins are considered to be involved in early onset of Alzheimer's disease. Presenilin 2 mainly localizes in large neurons. It exists in a heterodimeric complex formed from the endoproteolytically processed amino-terminal (34 kDa) and carboxy-terminal (about 23 kDa) fragments. This antibody is specific to presenilin 2. It will not bind presenilin 1.

Notable Publications

Author	Pubmed ID	Journal	Application
Jinghua Jin	36683147	Gut Microbes	WB
Bhupender Sharma	38595288	Neural Regen Res	WB

Storage


Storage:

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 16168-1-AP (PSEN2-Specific antibody) at dilution of 1:800 incubated at room temperature for 1.5 hours.



WB result of PSEN2-Specific antibody (16168-1-AP; 1:1200; incubated at room temperature for 1.5 hours) with sh-Control and sh-PSEN2-Specific transfected A549 cells.



Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 16168-1-AP (PSEN2-Specific Antibody) at dilution of 1:200 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 16168-1-AP (PSEN2-Specific Antibody) at dilution of 1:200 (under 40x lens).