

For Research Use Only

ACAT1 Polyclonal antibody, PBS Only

Catalog Number: 16215-1-PBS

Featured Product



Basic Information

Catalog Number:

16215-1-PBS

Size:

1 mg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG9215

GenBank Accession Number:

BC010942

GeneID (NCBI):

38

UNIPROT ID:

P24752

Full Name:

acetyl-Coenzyme A acetyltransferase

1

Calculated MW:

45 kDa

Observed MW:

38-45 kDa

Purification Method:

Antigen affinity purification

Applications

Tested Applications:

WB, IHC, IF/ICC, FC (Intra), IP, Indirect ELISA

Species Specificity:

human, mouse, rat

Background Information

ACAT1 (acetyl-CoA acetyltransferase), is a mitochondrial enzyme catalyzing the reversible formation or breakdown of acetoacetyl-CoA and plays important roles in ketogenesis and ketolysis. Deficiency of ACAT1 causes an inherited metabolic disorder. Recently ACAT1 has been reported to get involved in cancer progression.

Storage

Storage:

Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:

PBS Only

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com

W: ptgcn.com


This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data




WB result of ACAT1 antibody (16215-1-AP; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-ACAT1 transfected MCF-7 cells. This data was developed using the same antibody clone with 16215-1-PBS in a different storage buffer formulation.


Various lysates were subjected to SDS PAGE followed by western blot with 16215-1-AP (ACAT1 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 16215-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded human liver tissue slide using 16215-1-AP (ACAT1 antibody) at dilution of 1:100 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 16215-1-PBS in a different storage buffer formulation.




1X10⁶ HeLa cells were intracellularly stained with 0.4 ug Anti-Human ACAT1 (16215-1-AP) and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Isotype Control. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 16215-1-PBS in a different storage buffer formulation.




IP result of anti-ACAT1 (IP:16215-1-AP, 4ug; Detection:16215-1-AP 1:500) with mouse heart tissue lysate 4400ug. This data was developed using the same antibody clone with 16215-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded human liver using 16215-1-AP (ACAT1 antibody) at dilution of 1:100 (under 10x lens). This data was developed using the same antibody clone with 16215-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded human colon slide using 16215-1-AP (ACAT1 antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 16215-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using ACAT1 antibody (16215-1-AP) at dilution of 1:200 and Multi-rAb CoraLite® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002), CL594-Phalloidin (red). This data was developed using the same antibody clone with 16215-1-PBS in a different storage buffer formulation.